

Miocene vertebrates from Entre Ríos province, eastern Argentina

Alberto Luis CIONE¹, María de las Mercedes AZPELICUETA², Mariano BOND¹, Alfredo A. CARLINI¹, Jorge R. CASCIOTTA², Mario Alberto COZZUOL³, Marcelo de la FUENTE¹, Zulma GASPARINI¹, Francisco J. GOIN¹, Jorge NORIEGA⁴, Gustavo J. SCILLATOYANÉ¹, Leopoldo SOIBELZON¹, Eduardo Pedro TONNI¹, Diego VERZI², María Guiomar VUCETICH¹

Resumen.- VERTEBRADOS DEL MIOCENO DE LA PROVINCIA DE ENTRE RÍOS, ARGENTINA. La diversa fauna de antiguos vertebrados que se registra en los acantilados que bordean la margen oriental del río Paraná cerca de la ciudad de Paraná, provincia de Entre Ríos, Argentina se conoce científicamente desde la primera mitad del siglo XIX. En esos sedimentos se han colectado numerosos vertebrados de agua dulce, marinos y terrestres. Los fósiles proceden casi exclusivamente de la Formación Paraná (taxones marinos y de agua dulce: elasmobranquios, teleosteos, cetáceos, sirenios y pinnípedos) y en el "Conglomerado osífero" ("Mesopotamiense" auctorum) en la base de la Formación Ituzaingó (taxones marinos, de agua dulce y terrestres: elasmobranquios, teleosteos, cocodrilos, quelonios, aves y diferentes grupos de mamíferos). Los cetáceos sugieren que al menos el tope de la Formación Paraná es Tortoniano (Mioceno Tardío). El término "Piso Mesopotamiense" o "Mesopotamiense" es considerado inválido. El "Conglomerado osífero" parece representar un corto lapso. La fauna terrestre sugiere una edad Huayqueriense (Tortoniano) para el "Conglomerado osífero". La evidencia de vertebrados y las relaciones estratigráficas confirman la correlación de al menos la base de las capas puelchenses del subsuelo de la región pampeana con la Formación Ituzaingó. De acuerdo a la evidencia que aportan los cetáceos y los peces, las temperaturas marinas durante la depositación de la parte superior de la Formación Paraná eran similares a aquellas presentes en la plataforma atlántica actual a la misma latitud. Tanto la fauna terrestre como la de agua dulce del "Conglomerado osífero" indican un clima más cálido que el actual. Los vertebrados de agua dulce sugieren importantes conexiones entre las cuencas hidrográficas del sur y del norte de América del Sur. Los restos de aves y de mamíferos (y plantas) sugieren la presencia de áreas forestadas a lo largo de las costas de los ríos donde se depositó el "Conglomerado osífero" y áreas abiertas cercanas.

Introduction

The fossiliferous beds in the Paraná eastern riverside cliffs near the city of Paraná, eastern Argentina have been scientifically known since 1827 when Alcide D'Orbigny visited the area (D'Orbigny, 1842; Figures 1,2). The cliffs are extremely rich in marine and continental aquatic and terrestrial vertebrate remains. However, most of the collections were done without accurate stratigraphic provenance. For this, during many years it was debated if the material came from

¹ Departamento Científico Paleontología Vertebrados, Facultad de Ciencias Naturales y Museo, 1900 La Plata, Argentina. E-mail: acione@museo.fcnym.unlp.edu.ar, azpeli@museo.fcnym.unlp.edu.ar, palvert@museo.fcnym.unlp.edu.ar (Bond), acarlini@museo.fcnym.unlp.edu.ar, jrcas@museo.fcnym.unlp.edu.ar, mdlafu@museo.fcnym.unlp.edu.ar, zgaspari@museo.fcnym.unlp.edu.ar, fgoin@museo.fcnym.unlp.edu.ar, palvert@museo.fcnym.unlp.edu.ar (Scillato), palvert@museo.fcnym.unlp.edu.ar (Soibelzon), eptonni@museo.fcnym.unlp.edu.ar, dverzi@museo.fcnym.unlp.edu.ar, vucetich@museo.fcnym.unlp.edu.ar.

² Departamento Científico Zoología Vertebrados, Facultad de Ciencias Naturales y Museo, 1900 La Plata, Argentina. E-mail: dverzi@museo.fcnym.unlp.edu.ar.

³ Universidad Federal de Rondônia, Porto Velho, RO, Brasil. E-mail: cozzuol@enter-net.com.br

⁴ Centro de Investigaciones Científicas y Transferencia de Diamante, Dr. Materi y España 3105 Diamante, Argentina. E-mail: cidnoriega@infoshopdte.com.ar.

one level or another. Moreover, different marine levels were identified in the cliffs.

Presently, several of us are making new collections with accurate stratigraphic provenance in the cliffs between Pueblo Brugo and Diamante (ALC, JN, MMA, JRC, EPT) with financial support of the CONICET and ANPCYT. We are confident that many of the present uncertainties will be set with this field work.

In this paper, the vertebrate record from the marine and continental Miocene units in the Paraná area is briefly revised by different specialists and a preliminary revised list is reported. Finally the correlation of the bearing units, and climatic and biogeographic aspects are discussed.

We include great part of the literature dealing with Miocene vertebrates from Paraná area. The taxonomic lists include those taxa that are considered relatively well identified. Certainly, the investigations are in progress and, in the following years, many of the taxonomic assignments will be modified and new taxa will be included in the lists by means of new findings or the study of previous collections.

History

When Florentino Ameghino, in his important work on the extinct mammals of Argentina (1889), discussed the vertebrates from the «Piso Mesopotámico» of Doering (1882), he included about 90 mammal species in addition to fish and reptiles. This paper by Ameghino was a peak in the knowledge of the Miocene vertebrates from the Paraná area, which had begun with the visit of D'Orbigny to the area in 1827 (Figure 1).

The French researcher Alcide D'Orbigny arrived in America in 1826. In the next year, he explored the Paraná riverside cliffs near the city of the «Bajada de Santa Fé», a former name for the city of Paraná, and later (1842) described the stratigraphy and many fossils. One of these fossils was a species of the endemic ungulate genus *Toxodon* described by Laurillard in D'Orbigny (1842); the species was different from the more recent *Toxodon platensis*, described by Owen and coming from the «arcilla pampeana.» D'Orbigny, based on the invertebrates, correctly attributed a Tertiary age to the sediments.

Another famous scientist, Charles Darwin, visited the area during October, 1833. Darwin described the stratigraphy of the cliffs near Paraná. In a bed near the base of the ravines, he found shark teeth and invertebrates pertaining to extinct species. Above this level, Darwin found a consolidated bed that passed upward to typical Pampean sediments. Darwin interpreted that as the result of a large marine encroachment which was gradually transformed in a muddy estuary where corpses floated (Darwin, 1839).

The General Justo José de Urquiza, first president of the Confederación Argentina, with capital in the city of Paraná, propitiated the beginning of systematic geological and paleontological research in the riverside cliffs. Urquiza hired, among others, the belge Alfred Du Gratty, who in 1854 was director of the Museo de la Confederación in Paraná; the French physicist and naturalist Martin de Moussy, who was in charge of surveying the national territory; and the French architect (and amateur paleontologist), August Bravard, whose first assignment was surveying mineral deposits.

Bravard was appointed director of the Museo de la Confederación in 1857. A year later, the Registro Oficial de la Confederación published the «Monografía de los terrenos marinos terciarios de las cercanías de Paraná» (Bravard, 1858). This paper was the sum of his research which had previously been communicated in the natural history societies of Buenos Aires and Paris in 1855 and 1856, respectively. Bravard's paper agreed in several respects with the observations




FIGURE 1. Location map.

of D'Orbigny and Darwin but was in contradiction with the conclusions of a paper by De Moussy (1857).

Bravard (1858) described for the first time several teleosts and elasmobranchs, reptiles (including chelonians, ophidians and crocodiles), and mammals. The president Domingo Faustino Sarmiento sent the materials collected by Bravard to the Museo Nacional de Historia Natural of Buenos Aires. Mammals were represented by a whale and a rodent smaller than «*Megamys patagoniensis*,» described by D'Orbigny (1842) from the río Negro riverside cliffs near Carmen de Patagones (northern Patagonia). Bravard considered that the Paraná and Patagones beds were of similar age.

In 1861, Bravard died during an earthquake that destroyed the city of Mendoza (western Argentina). After Bravard (1858), the only publication in many years was the «Description Physique, Géographique et Statistique de la Confederation Argentine» (De Moussy, 1860).

The Museo de la Confederación was refounded as the Museo de la Provincia de Entre Ríos in 1884 and the teacher Pedro Scalabrini was its director since 1886. Scalabrini was a great fossil collector and soon was in contact with the great paleontologist Florentino Ameghino who eventually will describe the new material. This association was the origin of the knowledge of the mammal fauna from the «Piso Mesopotamiense» (Ameghino, 1883a, 1883b, 1885, 1886, 1889).

Between 1890 and 1892, the Swiss Kaspar Jacob Roth (Santiago Roth in the Argentinian literature; see Bond, 1999), surveyed the provinces of Entre Ríos and Corrientes. Roth made an important vertebrate collection from the «Conglomerado Osífero» of the Ituzaingó Formation, part of which was deposited in the Museo de La Plata, where he was appointed Head of the Sección Paleontología in 1895.

During the last part of the XIX and the beginning of the XX centuries, many amateurs collected near Paraná. Later, most of these private collections were deposited in the Museo Nacional de Historia Natural of Buenos Aires (Lelong Thévenet, Carles, Sors Cirera and Caixo collections) and Museo de La Plata (Sors Cirera). Other institutions where collections from Paraná area can be found are: Muséum National d'Histoire Naturelle of Paris and Natural History Museum of London.

During the XX century, since the important paper by Frenguelli (1920a), many authors published contributions on the paleontology and geology of the Paraná area (eg Bianchini and Bianchini, 1971; Aceñolaza, 1976; Aceñolaza and Sayago, 1980; Iriondo and Rodríguez, 1973; Noriega, 1995 and the bibliography cited therein).

Stratigraphic provenance of the Miocene vertebrates

The complex relationships between the marine and continental units cropping out in the cliffs provoked different interpretations. Most authors identified only one marine unit recognizable at the base of the cliffs (eg Ameghino, 1906; Scartascini, 1954; Aceñolaza, 1976). This marine unit is overlain by a thick fluvial and a terrestrial sequence. However, other authors proposed two or three marine transgressions (Frenguelli, 1920a; Cordini, 1949). The current stratigraphic scheme was proposed by Aceñolaza (1976; see also Aceñolaza and Aceñolaza, 1999; Figure 2) who interpreted the marine rocks as having originated during a single ingression represented by the Paraná Formation. Aceñolaza (1976) suggested that the marine beds located at higher levels in the riverside cliffs are relicts of an ancient topography that was wrongly




FIGURE 2. Generalized stratigraphic section of the late Cenozoic strata at the left side cliffs along the río Paraná near the city of Paraná, Entre Ríos, Argentina (modified from Aceñolaza, 1976). The arrow indicates the "Conglomerado osífero" at the base of Ituzaiño Formation.

interpreted as different marine incursions, especially by Frenguelli (1920a). The Paraná Formation is mainly composed by green mudstones and sandstones with oyster banks (Aceñolaza, 1976; Chebli et al., 1989). The Paraná Formation was apparently deposited during the large marine encroachment that covered the Chacopampean region during the middle Miocene ("Mid Transgressive Onlap Sequence;" see Uliana and Biddle, 1988; del Río, 1991). This transgression probably persisted in the East of Argentina until the Tortonian (see below). The fluvial Ituzaingó Formation overlies the marine unit. This formation is composed of a basal conglomerate ("Conglomerado osífero") with abundant vertebrate remains which is overlain by almost unfossiliferous whitish to yellow brown sandstones and green mudstones. Terrigenous Pleistocene (Ensenadan to Lujanian in the southern South American continental scale; see Cione and Tonni, 1995, 1996; Figure 3), poorly fossiliferous beds assigned to different units overlay the Ituzaingó Formation (Aceñolaza, 1976; Iriondo, 1980; Chebli et al., 1989). The Ituzaingó Formation (as Entre Ríos Formation) was correlated with the "Formación Puelche" of the Buenos Aires province subsoil (Reig, 1957; see below). According to the mammals occurring in the "Conglomerado osífero" and the stratigraphic relationships, the age of the base of Ituzaingó Formation is almost exclusively Tortonian (late Miocene) or Huayquerian in the local chronology (Pascual and Odreman Rivas, 1971; Cione, 1988; Figure 3; see discussion below).

The term "Piso Mesopotamiense" or "Mesopotamiense" was widely in use in the Argentinian vertebrate paleontology literature. This term was used for the first time by Doering (1882; see also Ameghino, 1883). Frenguelli (1920a) restricted the "Mesopotamiense" to the "Conglomerado osífero." Two of us discussed extensively the use of state/age concept in South America (Cione and Tonni, 1995, 1996) although we did not address the "Mesopotamiense" problem. Cozzuol (1993) proposed to define the "Mesopotamiense" as a formal stage/age unit.

Unfortunately, most of the fossils in collections from the Paraná area do not include adequate provenance. Labels in collections usually only state that the material come from the base of the cliffs near Paraná. However, after several campaigns, we confirmed that almost all the Miocene terrestrial and freshwater aquatic vertebrates come from the «Conglomerado Osífero» at the base of the Ituzaingó Formation. Field work is presently in progress and more precise information will be obtained. The «Conglomerado osífero» occurs in paleochannels, is laterally interrupted and rarely crops out because landslides. We do not find adequate to sustain a chronostratigraphic, biostratigraphic, geochronologic or even biochronologic unit based on the fossil and stratigraphic representation of the «Conglomerado osífero.» Consequently, we consider here the «Piso Mesopotamiense» sensu Frenguelli (1920a) or «Mesopotamiense» as invalid. In this paper, we will refer to the fossil content of the base («Conglomerado osífero») of a lithostratigraphic unit (Ituzaingó Formation) as present in several localities near Paraná without recognizing a local chronostratigraphic or biostratigraphic unit (Figure 2).

Some marine vertebrates (mainly sharks and rays) occur in the «Conglomerado osífero.» It has to be determined if the marine vertebrate remains were reworked from the Paraná Formation or they actually inhabited the channels where the «Conglomerado osífero» was deposited. Until now, we have corroborated that the Paraná Formation includes mainly marine, some freshwater and no terrestrial vertebrate.

In sum, present evidence indicates that most of the Miocene material from the Paraná area come from the marine Paraná Formation (elasmobranchs, teleosteans, cetaceans, sirenians and pinnipeds) and the «Conglomerado Osífero» at the base of the continental Ituzaingó Formation (elasmobranchs, teleosteans, crocodylians, chelonians, birds, and different groups of terrestrial and aquatic mammals).




FIGURE 3a y b. South American chronostratigraphic late Cenozoic continental units (modified from Cione and Tonni, 1995, 1996; Tonni et al., 1998) in comparison with the magnetostratigraphic scale (Cande and Kent, 1995) and the Global Stratigraphic Chart.

The marine vertebrate record

Chondrichthyans

(Alberto Luis Cione)

The marine elasmobranch fauna found in the Paraná Formation is composed by the odontaspimid *Carcharias acutissima* (abundant), the carcharhinids *Carcharhinus* spp. (abundant) and *Galeocerdo aduncus*, the heterodontid *Heterodontus*, the squatinid *Squatina*, the lamnid *Isurus hastalis*, the "otodontid" *Carcharocles megalodon*, the hemigaleid *Hemipristis serra*, the squalid *Squalus*, and batoids Dasyatidae and Myliobatoidei (D'Alessandri, 1896; Woodward, 1900; Priem, 1911; Frenguelli, 1920a, 1922; Cione, 1978, 1988; Arratia and Cione, 1996; Figure 4). The ichthyofauna has not been recently studied, but until now, there are no living species identified. Several of these species also occur (surely reworked in part) in the "Conglomerado osifero."

"Osteichthyans"

(Alberto Luis Cione, María de las Mercedes Azpelicueta and Jorge Casciotta)

Some marine fishes were described from indeterminate beds near Paraná. The sparid *Chrysophrys* sp., Sparidae? indet., and the labrid *Protautoga longidens* Alessandri, 1896 were mentioned (Alessandri, 1896). The material was lost in the First World War in Torino, Italy. However, from the illustrations it can be observed that the presence of Sparidae and Labridae is not supported. The material identified as Sparidae? indet. (Alessandri, 1896) is quite similar to a lower pharyngeal plate of the sciaenid *Pogonias cromis*. We have found sciaenid remains both in the Paraná Formation and the "Conglomerado osifero" (Cione and Torno, 1984; Figure 5).

Cetaceans

(Mario Alberto Cozzuol)

The family Pontoporiidae, represented here by *Pontistes rectifrons* Bravard, 1858, and an indeterminate smaller species (Cozzuol, 1985, 1993, 1996) was a group diverse and widely distributed in the Americas both in the Atlantic and Pacific during the Cenozoic. Presently, only one species, *Pontoporia blainvillei*, which is an inhabitant of the Argentinian biogeographic Province in the Atlantic coast of South America, is extant (Ridgway and Harrison, 1989).

Physeterid and balaenid remains are not determinable at lower taxonomic levels, but are similar to that of the recent species (Cozzuol, 1993, 1996).

Balaenoteridae are known by quite complete but undescribed cranial material, tentatively referred to the genus *Balaenoptera* (Cozzuol, 1993; 1996)

Sirenians

(Mario Alberto Cozzuol)

A tooth formerly assigned to the genus *Metaxyterium* (see Reinhart, 1976; Cozzuol, 1993) is presently considered as pertaining, with doubts, to genus *Dioplotherium* (see Cozzuol, 1996). Remains of pachiostotic ribs are also relatively common in the collections. These bones are characteristic of the sirenians, specially of the paraphyletic family Dugongidae.

Aquatic carnivores

(Mario Alberto Cozzuol)

The phocid *Properiptychus argentinus* is one of the two fossil pinnipeds known from Argentina.




FIGURE 3b




FIGURE 4. Neoselachians from the Paraná Formation. A-E, *Carcharias acutissima*; F, *Galeocerdo aduncus*; G, *Hemipristis serra*; H, *Carcharodes megalodon*; I-K, *Isurus hastalis*; Modified from Woodward (1900). All the material at natural size excepting H that is 2/3.

It was assigned to the paraphyletic subfamily «Monachinae» (see Muizon and Bond, 1982) and it was considered as primitive for the family, perhaps related to the paraphyletic genus *Monachus* (monk seals).

The continental vertebrate record

Chondrichthyans

(Alberto Luis Cione)

The genus *Potamotrygon* of the freshwater family Potamotrygonidae was mentioned by

Ameghino (1898), without figuration or description. Recently, Deynat and Brito (1994) assigned the large dermal scutes described by Larrazet (1886) from Paraná to the family Potamotrygonidae. Dermal scutes are found both in the marine Paraná Formation and the “Conglomerado osífero.” Part of the material probably correspond to other families.

«Osteichthyans»

(Alberto Luis Cione, María de las Mercedes Azpelicueta and Jorge Casciotta)

Numerous specimens of catfishes and characiforms, taxa that include the most conspicuous Neotropical freshwater fishes were found in the Ituzaingó Formation (and also in the Paraná Formation; Figure 5). Fishes from the Paraná area have been known since last century (Bravard, 1858). However, they have not been adequately studied yet (Ameghino, 1898; Woodward, 1900; Priem, 1911; see Cione, 1978, 1986; Arratia and Cione, 1996).

The catfish *Silurus agassizi* Bravard, 1858 was not described nor figured and must be considered *nomen nudum* (Cione, 1986). The assignation of a catfish skull to *S. agassizi* by Frenguelli (1920a) is obviously invalid. Catfishes of different families are present in the material from the “Conglomerado osífero” (ALC, personal observation). Some characiforms such as the serrasalmid *Colossoma macropomus* occur in the Ituzaingó Formation (Cione, 1986; Figure 5). This species has been also identified in La Venta Group (Middle Miocene of Colombia; Lundberg et al., 1986, 1988). The records from the Ituzaingó Formation and La Venta Group corroborate the




FIGURE 5. Teleosteans from the Ituzaingó Formation. A, *Colossoma macropomum*, right dentary, anterior view. B, Cynodontidae indet., tooth, lateral view (modified from Cione and Casciotta, 1997). C, cf. Sciaenidae, right dentary, labial view. The lines are 1 cm.

biogeographic relationship based on by other faunal elements (eg. crocodylians, trichechids, cetaceans).

Canine teeth of the characiform family Cynodontidae, similar to genera *Hydrolicus*, *Cynodon* and *Raphiodon*, were described from the “Conglomerado Osífero” (Cione and Casciotta, 1997; Figure 5). However, the size and position of serrae on cuttig edges is different from that recorded in species of these extant genera. One large tooth probably corresponds to a specimen of more than 1000 mm SL.

A tooth identified as pertaining to the ginglymodi *Lepisosteus* (Alessandri, 1896) is assignable to an indeterminate Crocodylia (Cione, 1986).

“Reptilians”

(Zulma Gasparini and Marcelo de la Fuente)

The first references of fossil crocodiles, turtles and ophidians of Argentina, are those of Bravard (1858). Further papers, describing fragmentary material from the “Conglomerado osífero,” supported the determination of numerous species of chelids, testudinids, alligatorids, gavialids, and teiids (Burmeister, 1885; Ambrosetti, 1890, 1893; Wieland, 1923; Rusconi, 1933, 1934, 1935; Mones, 1986). However, this high diversity was reduced by more recent studies (Báez and Gasparini, 1977; Donadío, 1983; Gasparini et al., 1986, de la Fuente, 1988, 1992; Broin and de la Fuente, 1993; Albino, 1996; Gasparini, 1996; Langston and Gasparini, 1997).

A single chelid (*Phrynops* cf. *Phrynops geoffroanus* complex) (see Rhodin and Mittermeier, 1983; Figure 6), one indeterminate testudinid, three species of alligatorids (*Caiman jacare*, *C. latirostris* and probably *C. australis*; Figure 6), one nettosuchid (*Mourasuchus nativus*), one gavialid (*Gryposuchus neogaeus*; Figure 6), and one teiid (*Tupinambis* cf. *T. meriana*) are presently accepted.

The diversity of crocodiles in the area of Paraná may be compared to that recorded currently in the basins of the large northern South American rivers, such as Paraguaí and Solimões rivers in Brazil and Madre de Dios river in Perú (Medem, 1983). However, it is richer at the familial level, as there are gavialids (currently limited to southeastern Asia) and netosuchids (extinct). Likewise, *Caiman jacare* and *Caiman latirostris* were recorded in “Conglomerado osífero;” both species inhabiting nowadays the Mesopotamian margin of the Paraná river basin.

Chelonian diversity is lower at the familial and generic level than that of other coeval South American sites. For example, podocnemidids are not represented in the Ituzaingó Formation; among chelids, no species referable to *Chelus* has been recorded, and the diversity of testudinines is much lower (cf Lapparent de Broin et al., 1993, and literature therein; Wood, 1976, 1997; Wood and Díaz de Gamero, 1971). Concerning chelid turtles, the greatest generic and even specific affinities is found to chelids of the Kiyú Formation, Department of San José, Uruguay (cf Perea et al., 1996; their Figure 2).

Birds

(Jorge Noriega)

The bird record from the «Conglomerado osífero» comprises eight species of six orders: Pelecaniformes, Charadriiformes, Anseriformes, Ciconiiformes, Rheiformes and Gruiformes.

The Pelecaniformes are represented by the largest known darter (*Macranhinga paranensis*, Anhingidae; Figure 6). This species showed mechanisms of aquatic and aerial locomotion similar to those of the recent putative sister group of the Anhingidae, i.e. the family Phalacrocoracidae (Noriega, in press).

Another indeterminate darter seems to have been a flightless species, with a size similar to




FIGURE 6. “Reptiles” and bird from the Ituzaingó Formation. *Phrynops* cf. *P. goffroanus*, caparace: A, dorsal view; B, visceral view (modified from Wieland, 1923); the line is 10 cm. C, *Gryphosuchus neogaeus*, skull, dorsal view (modified from Gasparini, 1968); the line is 10 cm. D, *Caiman latirostris*, skull, ventral view (modified from Gasparini, 1968); the line is 2 cm. E, *Macranhinga paranensis*, tarsometatarsus, anterior view (modified from Noriega, 1992); the line is 1 cm.

the recent species *Anhinga anhinga* (Noriega, 1994, 1995).

The Charadriiformes record include an indeterminate species of genus *Megapaloelodus* from the extinct flamingo family Palaeolidae and an indeterminate Phoenicopteridae (Noriega, 1995).

The only fossil South American representant of the subfamily Dentrecheninae (Anseriformes, Anatidae) occurs in the «Conglomerado osifero.» This subfamily is more related to the Dendrocygninae (whistling ducks) and Anserinae (geese and swans) than to the real ducks (Anatinae; Noriega, 1995).

The ciconiiform Mycteriini storks are only known in the Tertiary of South America in the «Conglomerado osifero» (Noriega, 1994, 1995).

Cursorial birds are represented by indeterminate reiform Rheidae, gruiform Rallidae and Phororhacidae (Noriega, 1995). This last family includes the large phororhacine *Onactornis?* and the medium sized palaeociconine *Andalgalornis*. Phororhacid taxa from the «Conglomerado osifero» must be revised (Noriega, 2000).

Marsupials

(Francisco Goin)

Three carnivorous marsupials (Sparassodonta), and three “opossum-like” marsupials (Didelphimorphia) are known from the “Conglomerado osifero:” *Notictis ortizi* Ameghino 1889 (Sparassodonta, Hathliacynidae) is a very small hathliacynid solely represented by a partially preserved dentary. *Stylocynus paranensis* Mercerat 1917 (Sparassodonta, Borhyaenidae, Prothylacyninae) is one of the largest known prothylacynine marsupials. Its mandibular and dental specializations suggest a predominantly omnivorous diet (Marshall, 1979). *Achlysictis lelongi* Ameghino 1891 (Sparassodonta, Thylacosmilidae), one of the most spectacular, large predaceous marsupials evolved in South America, is also present at Paraná deposits. Goin and Pascual (1987) proposed to keep the name *Thylacosmilus atrox* to this and all other late Miocene-Pliocene known thylacosmilids (but see Marshall et al., 1990). *Philander entrerrianus* Ameghino 1899 (Didelphimorphia, Didelphidae, Didelphinae) is probably a synonym of the Recent *Philander opossum*, and its stratigraphic provenance (“Conglomerado osifero”) should be regarded as highly dubious (Goin, 1997b; see below). *Chironectes* sp. (Didelphimorphia, Didelphidae, Didelphinae), also recorded from “Barrancas del Paraná, Mesopotamiense” (original label) is also a probable synonym of the Recent species *Chironectes minimus* (Reig, 1958; Reig et al., 1987; Marshall, 1987; but see Goin, 1991). It is another case of a possible Recent opossum mixed among “Mesopotamian” taxa. *Zygolestes paranensis* Ameghino 1898 (Didelphimorphia, Didelphidae, Marmosinae) is a curious, tiny marmosine opossum whose affinities were unclear until very recent times. On account of its peculiar features noted by Reig (1957) in his diagnosis of the species, Reig et al. (1985, 1987) regarded it as a Didelphoidea incertae sedis. Marshall (1987) and Marshall et al. (1990) recognized for it a new Tribe among the Didelphinae: Zygolestini. Goin (1991, 1995, 1997b) and Goin et al. (MS) argued in favour of its Marmosini affinities, especially with species of *Graulinanus*. A recently described new species of *Zygolestes*, from Huayquerian levels of Pampean area, suggests to Goin et al. (in preparation) that a Miocene age for *Zygolestes paranensis* is reasonable.

Xenarthrans (Figure 7)

(Gustavo J. Scillato Yané and Alfredo A. Carlini)

Except for a few prior findings of vertebrates that have only historical interest, the discovery




FIGURE 7. Xenarthans and litoptern from the Ituzaingó Formation. A, *Palaeohoplophorus scalabrini*, lateral view of causal tube. B, *Pseudoeuryurus lelongianus*, dermal scute of the dorsal caparace. C, *Scalabrinitherium bravardi*, upper molar. D, *Palaeohoplophorus scalabrini*, dermal scute of the dorsal caparace. E, *Kraglievichia paranensis*, dermal scutes in dorsal view. (A-D, modified from Ameghino, 1889; E, modified from Edmund, 1987). The line is 1 cm.

and study of the xenarthrans from the Ituzaingó Formation began with the collections of Pedro Scalabrini. He sent the remains to F. Ameghino for study (see Ameghino 1883, 1883a, 1885, 1886 and 1889). Later, numerous xenarthrans, mainly tardigrades, were re-studied or recognized as new ones by L. Kraglievich (1922, 1923, *passim*) and Scillato-Yané (1975a, 1980, 1981, 1982; Pascual *et al.* 1990). Recently, in relation with new collections, different remains of tardigrades were described from the Ituzaingó Formation and systematic and biostratigraphic problems were addressed (Carlini *et al.* in press). Likewise, a complete list of the xenarthran species of the Argentine Cenozoic distributed according to stratigraphic units and geographic areas is given elsewhere (Scillato Yané and Carlini, in press). In these analyses, the xenarthran diversity from the Ituzaingó Formation (42 genera and 56 species, see Table 1) is found to be exceeded only by that of the Santa Cruz Formation (Santacrucian, early Miocene from Patagonia), and much greater than that of any other South American unit. However, the nature of the deposits and the fragmentary and dissociated remains have led to both an overestimation of the diversity (by recognizing species based on non-homologous remains), and an underestimation when the assignment of specific differences was impossible in view of the fragmentary materials.

Among the Cingulata, dasypodids are very scarce in the "Conglomerado osífero," all the cited species being known by a single specimen, except for *Proeuphractus limpidus*. This is particularly outstanding, especially in comparison with the abundant record in the coeval sites of northwestern Argentina.

The record of *Zaedyx* is dubious in view of the poor material. Besides, the occurrence of this taxon disagrees with the remaining taxa, because it is typical from relatively cold and xeric regions.

Pamphateres were hitherto represented by a single but quite frequent species, *Kraglievichia paranense*. Recently, a new species of genus *Scirrotherium* was identified in the "Conglomerado osífero" and will be described in a future paper. The genus *Scirrotherium* occurs for the first time outside the middle Miocene of La Venta (Colombia).

Glyptodontids are very diversified in the Ituzaingó Formation (12 genera and 12 species). The tribe Palaehoplophorini is highly dominant in taxic composition which is in contrast with the "Araucanian" of northwestern Argentina, where they are absent, and the very scarce record in the Miocene-Pliocene of the Pampean region. The occurrence of *Berthawylexia* (Miocene-Pliocene from Uruguay and Pliocene of the Pampean region) is the first for the Ituzaingó Formation. "*Hoplophorus*" *verus* does not belong in this Pleistocenic genus, but perhaps in *Hoplophractus* Cabrera or *Eosclerocalyptus* C. Ameghino. The available material of *Trachycalyptus? angulatus* is not enough to determine whether it is assignable to this genus or to *Comaphoropus*.

The largest glyptodont of the Ituzaingó Formation is the neuryurini *Urotherium interundatum*, with a size similar to *Neuryurus rudis* (Pleistocene of the Pampean region). The doedicurine *Comaphorus concisus* is similar to *Urotherium interundatum* in the scute surface and the profile of the elevated central figure. Unfortunately, the material is not enough as to accurately establish the variability of the different carapace regions. However, carapace similarities clearly suggest a close relationship between Sclerocalyptinae Neuryurini and Doedicurinae (see also Ameghino 1889:840 ss.). *Pseudoeuryurus lelongianus* also has a similar scute morphology to *C. concisus* and *U. interundatum*.

The diversity of the tardigrades from the Ituzaingó Formation is higher than that of any other unit of the late Miocene-Pliocene of Argentina. The 23 genera and 37 species recognized are distributed among the families Megatheriidae (Megatheriinae and Nothrotheriinae), Megalonychidae (Orthotheriinae, Megalocninae and Megalonychinae) and Mylodontidae (Mylodontinae, Octomylodontinae and Scelidotheriinae). These families include few small species




FIGURE 8. Rodents from the Ituzaingó Formation. A, *Phoberomys* sp., lower right molar, occlusal view. B, *Phoberomys*, lower right premolar, occlusal view. C, *Eumegamys paranensis*, lower right molariform series. Modified from Kraglievich, 1932. The line is 1 cm.

but those middle and large sized are better represented.

Megatheriinae genera already cited in the Ituzaingó Formation are: 1- *Eomegatherium* (also in the Mayoan and in the upper Miocene of San Gregorio cliffs, Uruguay); 2- *Promegatherium* (probably also in the basal beds of the Huayquerías Formation, late Miocene of Mendoza; and 3- *Pliomegatherium*, the only exclusive to the “Conglomerado osífero.” These genera are represented by a smaller species with more primitive characters than those coeval from northwestern Argentina, the Huayquerías of Mendoza and northern Patagonia.

A right humerus, the measures of which reach those of *Megatherium americanum* is quite larger than the remaining megatherines and different in morphology to the other genera from the Ituzaingó Formation. However, when compared with remains of *Pyramiodontherium* from Catamarca (“Araucanian”), it seems to coincide in the diaphysis section and the progression of its proximal diameter. Consequently, we consider this fossil as pertaining to that genus (perhaps a new species; Carlini *et al.*, in preparation). *Pyramiodontherium*, with a single species (*P. bergi*) has been hitherto recognized exclusively in the “Araucanian” of Catamarca and Tucumán.

The nothrotherine are represented by *Neohapalops*, an endemite, and *Pronothrotherium*, which is also known in the Miocene-Pliocene of the Uruguay and northwestern Argentina and the Pliocene of Mendoza and the Pampean region.

The diversity of Megalonychidae from the Ituzaingó Formation is higher than that of any other South American unit and only can be compared with that of the Pleistocene of West Indies. The “Orthrotheriinae” (which probably is not a natural group, Carlini and Scillato-Yané,

in preparation) are the most abundant, represented by the genera *Pliomorphus*, *Menilaus?*, *Torcellia*, *Paranabradys* and *Orthotherium*. The five nominated species of *Orthotherium* must be revised. Among megalonychids there are genera related to North American forms (*Protomegalonyx*, Megalonychinae) and to West Indies megalocnines (*Amphiocnus*, see Pascual et al. 1990 and probably *Pliomorphus*, in study). *Megalonychops* is the largest known Megalonychidae in the Ituzaingó Formation, comparable in size to *Megatherium*. Remains of *Megalonychops*, which also occurs in the Miocene-Pliocene of Uruguay and the Pleistocene of the Pampean region, are scarce in the “Conglomerado osífero” and need to be restudied. *Diodomus copei* is probably a Megalonychidae but the scarce remains should be revised.

Remains of mylodontid subfamilies are quite irregular in diversity and frequency. Mylodontinae reach higher generic and specific diversity and abundance in the “Conglomerado osífero” than in any other Tertiary stratigraphic unit. They are abundant in the Miocene-Pliocene of northwestern Argentina but very scarce in coeval localities of the Pampean region.

The peculiar Octomyodontinae (see Scillato-Yané, 1977a) are only known in the “Conglomerado osífero” and the Vivero Member of the Arroyo Chasicó Formation in the Pampean region (represented by *Octomyodon robertoscagliai*).

The Scelidotheriinae are very scarce. We have seen molariforms of a rather large scelidotherine coming from the Ituzaingó Formation.

“Native ungulates” (Figure 7)

(Mariano Bond)

After the first descriptions by D'Orbigny (1842) and Bravard (1858), the systematic description of ungulates from Paraná begun in the decade of 1880 (Burmeister, 1885-1891 and especially Ameghino, 1883a,b, 1885, 1886, 1887, 1889, 1891, 1904a, b, 1906; Frenguelli, 1920b; Kraglievich, 1931, 1934). Pascual and Odreman (1971) apported a list and some comments. The study of protherotheriids by Bianchini and Bianchini (1971) is the only recent revision of native ungulates from the Ituzaingó Formation.

In the «Conglomerado osífero», the two orders that survived Chasicóan times are present: Litopterna (Macrauchiinae and Protherotheriidae) and Notoungulata.

Macrauchiinae are represented by the subfamily Macrauchiinae, which is recorded since the Chasicóan (late Miocene). All macrauchiinae recorded seem to be more generalized than *Promacrauchenia* (Montehermosan and Chapadmalalan) and more related to taxa from the Chasicóan and Huyquerian (Bond and López, 1996). The Protherotheriidae Protherotheriinae correspond to more generalized taxa than those from the Montehermosan while there persist some «Pansantacrucian» species. Comparing with other faunas from the late Miocene and Pliocene beds of Argentina, Bolivia and Uruguay, Litopterna are extremely abundant (as numerous as Notoungulata) in the «Conglomerado osífero.»

Notoungulata present differences in diversity and systematic composition in relation to other faunas of late Miocene and Pliocene units of Argentina, Bolivia and Uruguay. There are no «rodentiform» Tipotheriidae, Hegetotheriidae Hegetotheriinae and Pachyrhinae, nor Mesotheriidae Mesotheriinae while Hegetotheriidae, especially Pachyrhinae, and Mesotheriidae are extremely frequent in the other localities.

The record of Mesotheriidae (Ameghino, 1906) and Pachyrhinae (Pascual and Odreman Rivas, 1971) from the «Conglomerado osífero» seems to have been based on an erroneous identification (Kraglievich, 1931; Bond and López, 1998). However, in the «Conglomerado osífero» the last probable representants of the Intertheriidae Intertheriinae occur. *Protypotherium antiquum* is also the largest species of Intertheriinae, subfamily which is very frequent in the




FIGURE 9. Zoogeographic pattern of Argentina (modified from Ringelet, 1961). 1, Subtropical Dominion. 2, Pampean Dominion. 3, Andean Dominion. 4, Central or Subandean Dominion. 5, Austro-Cordilleran Dominion.

Santacrucian and Colloncuran of Patagonia. The Interatheriinae occurs in Chasican beds for the last time in the Pampas. Some probable Interatheriinae remains are assignable to the genus *Munizia*, an endemic and larger genus.

Toxodontids are the more abundant and diverse Notoungulata in the «Conglomerado osífero.» Different groups with euhypsodont molariforms recognized from the Colloncuran, Mayoan and Chasican are present. The Xotodontinae and Toxodontinae include medium and large sized taxa and the Haplodontheriinae and Dinotoxodontinae with large to gigantic sized taxa. The supposed occurrence of Nesodontinae (Ameghino, 1891, 1906) is probably based on remains of a more advanced toxodontid known from the Huayquerian of Uruguay (?*Berroia*, Kraglievich, 1934). Xotodontinae have the last certain record in the Montehermosan and probably in the Chapadmalalan. Haplodontheriinae, large to gigantic toxodontids, include several taxa with a remarkable frontal bony protuberance which would indicate the occurrence of a «horn.» This family of «rhinocerotid» notoungulates is recorded for the last time in the Montehermosan. Dinotoxodontinae, a group not clearly defined of large toxodontids characterized by a notable lower projection of the mandibular symphysis, seems to have been abundant in the Pampean area during the Chasican and Huayquerian, and presents several forms in the «Conglomerado osífero.» The abundance of toxodontids appears to agree with the traditional attribution of semiaquatic habits, or at least relation with wet environments, of these notoungulates.

The toxodonts from the «Conglomerado osífero» have been described mainly on the basis of isolated teeth. However, they have been used for recognizing some subgroups inside the Family Toxodontidae (eg *Haplodontherium* and the Haplodontheriinae). Besides, it is evident that there is similitude between many taxa from the Ituzaingó Formation and those of the Miocene-Pliocene from Argentina, Bolivia and especially Uruguay and Acre in Brazil. For this, only a thorough revision of the taxa described for the Ituzaingó Formation will shed light not only on the systematic but on the relationships of these taxa with those from the other coeval units.

Rodents (Figure 8)

(María Guiomar Vucetich and Diego Verzi)

Hystricognathi rodents are represented by families Echimyidae, Myocastoridae, Caviidae, Hydrochoeridae, Dinomyidae, Neopiblemidae, Chichillidae, Abrocomidae and Erethizontidae. Probably there is no other so diverse (at familial level) rodent fauna in the Cenozoic of South America. Octodontidae is the only southern South American family that is not present. This familial diversity would indicate the influence of different biogeographic domains. For example, the presence of abrocomids and chinchillids suggests a connection with central and western Argentina while that of neopiblemids indicate a clear connection with the Miocene beds of the Brazilian Acre region.

The rodent fauna of the «Conglomerado osífero» is also very rich at the generic level, including more than 40 genera. The richest families are Dinomyidae (16), Hydrochoeridae (9), and Caviidae (6). The remaining families are represented by one to three genera (Table 3). Rodents from the «Conglomerado osífero» have not been revised exhaustively after the original descriptions. However, a preliminar revision of type and calcotypes indicates that some species are based on jaw fragments, or juveniles, and even on one isolated molariform. For this, it is probable that the number of species and genera could be overestimated.

Families Dinomyidae, Hydrochoeridae and Caviidae show their highest diversity in the «Conglomerado osífero», with several genera, some of which are endemites.

Recently, with the revision of the calcotype of *Paradoxomys cancrivorus*, a former enigmatic

taxon, it was demonstrated that it represents the first Erethizontidae for the “Conglomerado osifero.” *Paradoxomys cacrivorus* seems to be related to the radiation of recent taxa more than to the Patagonian radiation (Vucetich and Candela, in preparation).

The presence of *Protabrocoma*, related to the genus *Abrocoma* is remarkable because this latter is restricted to the arid Andean region. The most plausible hypothesis is that *Protabrocoma paranensis* had environmental constraints different to the living species of *Abrocoma*. In this sense, it is remarkable the record of a recent unusual abrocomid, showing climbing capabilities, from the forested areas in the northern Vilcabamba cordillera (Bolivia; *Cuscomys*, Emmons, 1999).

Echimyids are represented by endemic taxa of the Eumysopinae. These taxa would represent different lineages to the Eumysopinae of the Late Miocene of central and western Argentina (Verzi et al., 1995; Vucetich and Verzi, 1995) and the Pliocene and Pleistocene of the Pampean area (Vucetich and Verzi, 1996; Vucetich et al., 1997).

A characteristic of the “Conglomerado osifero” (and coeval localities) is the presence of the largest representants of the order Rodentia (Neopiblemidae, Hydrochoeridae and Dinomyidae).

Terrestrial carnivores

(Leopoldo Soibelzon)

The first procyonid carnivore recognized in Argentina is the procyonine *Cyonasua argentina* (see Ameghino 1885, 1886) from the “Conglomerado osifero.” Procyonid are medium-sized, good swimmers omnivores that mainly inhabit forested environments (Bond, 1986). Their habits probably facilitated the early entrance of these carnivores as “waif immigrants” from North America during the Huayquerian (late Miocene; Pascual et al., 1985).

Cetaceans

(Mario Alberto Cozzuol)

At least three species of the Family Iniidae and a Pontoporidae were described from the «Conglomerado osifero.» *Ischyorhynchus vanbenedeni* Ameghino, 1891 is the best known cetacean from the Paraná area (Cozzuol, 1993, 1996). It presents morphological adaptations for a turbid fluvial environment (Pilleri and Gühr, 1979).

Sauroctes argentinus Burmeister, 1871, also a relatively abundant species, is known by partial skulls and mandibles (Cozzuol, 1985).

Both species were also recorded in the Miocene beds of Acre and Uruguay (Boquentin et al., 1989; Rancy et al., 1989; Perea and Verde, 1982; personal observation, MAC).

Other species of *Sauroctes*, *S. gigas* (Cozzuol, 1989) is remarkable for its unusual size for a freshwater cetacean (about 4.5-5 m).

Sirenians

(Mario Alberto Cozzuol)

The occurrence of a Trichechidae (*Ribodon limbatus* Ameghino, 1883) at this latitude is important from a biogeographic and climatic perspective. But *R. limbatus* is also remarkable because this species shows a type of tooth change which is considered advanced in comparison with primitive trichechids, such as *Potamosiren* (from Laventan beds of Colombia) and considered a synapomorphy shared with the recent *Trichechus* (Domning, 1982; Cozzuol, 1993, 1996).

R. limbatus was also recorded in the Acre region, at the Peruvian-Brazilian border (Frailey, 1986; Cozzuol, 1993; 1996). A maxilar of a sirenian from the coastal plain of South Carolina, USA, was referred as belonging to an unidentified species of *Ribodon* (Domning, 1982). The age




FIGURE 10. Ichthyogeographic «provinces» of southern South America according to Ringuélet (1975), modified by Arratia et al (1983). A1-A3: Austral Subregion. A1, Chilean Province; A2, Cuyan Province; A3, Patagonic Province. B1-B18: BRazilian Subregion; B1, Titicaca Province; A3 Patagonic Province. B1-B18: Brazilian Subregion; B1 Titicaca Province; B2-B4, Paranean Domain; B2, High Paraguay Province; B3, High Paraná Province; B4, Pano-Platan Province; B5, coastal river of South eastern Brazil Province..

of the fossil was mentioned as probably early Pliocene. Taking into account those data, it seems that *Ribodon* had a similar geographic range than the living *Trichechus*, except by the African coast.

Stratigraphic correlation

Marine

Sharks are suggestive of Middle Miocene to Pliocene age but cetaceans suggest, more accurately, a Late Miocene age for the top of the Paraná Formation. The occurrence of Balaenidae and Physeteridae similar to modern taxa, of Balaenopteridae (unknown in beds older than late Miocene), and the lack of common taxa with the early Miocene Gaiman Formation (Cione and Cozzuol, in preparation) and the Middle Miocene Puerto Madryn Formation (Cozzuol, 1993, 1996), suggests that the uppermost part of the Paraná Formation (at least in the outcrops in the bottom of the Arroyo Ensenada valley and Villa Urquiza) would be Tortonian in age. Besides, there are no outcrops with typically Chasicosan fauna (early Tortonian in age) overlying the Paraná Formation and underlying the «Conglomerado osífero» of the Ituzaingó Formation, where Huayquerian fauna occurs (see below). Certainly, both formations are separated by an important unconformity and an important lapse is perhaps not represented by sediments (and fossils).

Continental

The exact age of the Ituzaingó Formation has been largely discussed. Mammals from the Ituzaingó Formation are extremely diverse. At first sight, the coexistence of so many taxa would be suggestive of an artificial mixture, especially considering that the ecological requirements of many of them must have been very similar and the methodology of collection of former researchers. Some authors even proposed that the mammal material from the “Conglomerado osífero” could be redeposited from beds of different age: Chasicosan, Huayquerian, Montehermosan and even Santacrucian (Bianchini and Bianchini, 1971; Scillato-Yané 1977a, 1980, 1981; Marshall et al., 1983). Another important drawback is the accurate identification of taxa because most of the determinations are old and should be revised. Consequently, we believe that considerably field work and taxonomic studies should still be done in all Late Miocene units in southern South America.

However, we consider that:

1. The age of the “Conglomerado osífero” is constrained by that of the underlying Paraná Formation (Figure 2).
2. According to cetacean evidence, the top of the Paraná Formation seems to be early Tortonian in age.
3. There is an important unconformity between the Paraná and Ituzaingó Formations that could represent a relatively long lapse.
4. All the terrestrial vertebrate remains are disarticulated. However, this is usual in a fluvial deposit. The only articulated vertebrate recorded are some freshwater fishes very well preserved into nodules. Remarkably, many bones are unworn, especially taking into account the conglomeradic and sandy composition of the deposit. We regard unlikely that the material would come from other terrestrial units (which actually are unknown in the area; see also Cozzuol, 1993). Moreover, the fluvial basin developed onto the marine beds and there is no doubt that some (not necessarily all) marine vertebrates remains present in the “Conglomerado osífero” were reworked from the Paraná Formation. We did not find terrestrial

vertebrates in the Paraná Formation so far.

5. The major affinities of the “Conglomerado osífero” are with Chasicosan and Huayquerian faunas (Figure 3; Tables 1-3). Genera shared (figure in parenthesis: certain records plus doubtful records): Chasicosan, 11 (17?); Huayquerian, 28(32?). We discard the correlation with Santacrucian, Mayoan, Laventan and Montehermosan. The occurrence of a few taxa with Santacrucian, Mayoan and Laventan affinities is here considered as the persistence of ancient lineages (see above). There are several other cases in the Cenozoic of South America. Affinities with Montehermosan beds are reduced to the first occurrence in both unit of the species *Macroeuphractus retusus* (three scutes, see Scillato Yané, 1975b) and *Protohydrocoerus* (a molar, MGV, personal observation). Consequently, the discussion is reduced to the correlation with Chasicosan or Huayquerian beds.
6. The Chasicosan age was recognized on the basis of the stratigraphic and faunistic record of the Arroyo Chasicó Formation in southern Pampean region (for a comment see Bondesio *et al.*, 1980; Marshall *et al.*, 1983; Tonni *et al.*, 1998). The Loro Huasi Formation (Valle de Santa María, Catamarca, Galván and Ruiz Huidobro, 1965; «Chiquimil» or «Chiquimil Formation» of other authors, see Marshall and Patterson, 1981) is perhaps correlated to the type Chasicosan (Marshall *et al.*, 1983).
7. The Huayquerian age is mainly recognized on the basis of the stratigraphic and faunistic record of the Andalhuala Formation (Valle de Santa María, Catamarca, Galván and Ruiz Huidobro, 1965; «Araucanense» of Frenguelli, 1930; Riggs and Patterson, 1939 non «Araucanense» of Ameghino, 1906 and other authors, a much more embracing concept) and the Epecuén «Formation» and Cerro Azul Formation (Pampean area; Pascual *et al.*, 1965; for other Huayquerian localities, see Marshall *et al.*, 1983). The Andalhuala Formation is represented by the levels XV to XX of Chiquimil area and 14 to 18 of Corral Quemado area, both in Catamarca province (see Stahelcker's stratigraphic sections in Riggs and Patterson, 1939 and Marshall and Patterson, 1981). We proposed elsewhere that the levels 18 to 21 of Corral Quemado might not be Montehermosan but Chapadmalalan in age as well as several sections in Bolivia (Cione and Tonni, 1996).
8. For analyzing the first mammal occurrences in the Ituzaingó Formation and other sequences we preferred to use the fossil record of the Chasicosan type section, the Huayquerian Andalhuala Formation and the Huayquerian beds of the Pampean region. The Andalhuala Formation section should be the actual stratotype of the Huayquerian. The Arroyo Chasicó Formation is well sampled and is overlain by Huayquerian beds (Tonni *et al.*, 1998). The thick sequences in the Catamarca province were well sampled, were radiometrically and magnetostratigraphically analyzed, and include a possible Chasicosan section (Loro Huasi Formation), a Huayquerian section (Andalhuala Formation), and a probable Chapadmalalan section. The Epecuén and Cerro Azul Formations do not crop out in a stratigraphic sequence with older and younger units but are rich in mammal content.
9. For correlation we considered relevant the first occurrences and the shared occurrence of several mammal taxa mostly at supraspecific level due to the lack of a recent systematic revision in most the groups.
10. The «Conglomerado osífero» shares first mammal occurrences with (Tables 1-3): 1) the Andalhuala Formation: *Myocastoridae*, *Parahoplophorus*, *Pyramiodontherium*, *Pronothrotherium*, *Sphenotherus*, *Neuryurini* (*Urotherium*) and *Xotodon*. 2) Huayquerian localities in Buenos Aires and La Pampa provinces: *Cavidon*, *Protabrocoma*, *Plohophorus?*, *Tetrastylomys*, and *Zygolestes*. 3) Both the Andalhuala Formation and Huayquerian localities in Buenos Aires and La Pampa provinces: *Achilisticis*, *Macroeuphractus*, *Doedicurinae* (*Eleutherocercus*), *Kiyutherium*, *Paleocavia*,

- Promacrauchenia?*, *Cyonasua* (*Cyonasua*), and *Achlysictis*. 4) The type Chasicoan: *Tetrastylus*, *Potamarchus*, *Carlesia*, *Diaphoromys*, *Gyriabrus*, *Lagostomopsis*, *Cardionmys*, *Parodymys?*, *Cardyatherium*, *Chasicotatus*, *Proeuphractus*, *Kraglievichia*, Octomyodontinae (*Octomyodon*), *Protomegalonyx*, *Brachytherium*, *Toxodotherium* (in Chiquimil), *Dinotoxodon*, *Protypotherium*, and *Cullinia?* Consequently, the «Conglomerado osífero» shares 18 (20?) first generic occurrences with Huayquerian localities and 16 (19?) with Chasicoan localities (figure in parenthesis: certain records plus doubtful records).
11. The «Conglomerado osífero» also shares with the type Chasicoan several taxa that are unknown in other levels (Tables 1-3): *Parodymys* (Vivero Member), *Octomyodon*, *Protomegalonyx*, *Protypotherium* (genus that is known from the Santacrucian) and *Cullinia?* However, *Protomegalonyx*, *Octomyodon* and *Protypotherium* correspond certainly or probably to different species (Scillato Yané, 1977a,b; MB, personal observation), *Parodymys* is based on one tooth and the identification of *Cullinia* in the «Conglomerado osífero» is uncertain (Vucetich and Bond, personal observations).
 12. The «Conglomerado osífero» shares with the Andalhuala Formation and the Pampean Huayquerian localities several taxa that are unknown from older levels: *Parahoplophorus*, *Pyramiodotherium*, *Sphenotherus*, *Protabrocoma*, *Tetrastylomys* and *Kiyutherium*. Besides, the «Conglomerado osífero» shares with the Andalhuala Formation or the Pampean Huayquerian localities several species that are unknown from older levels: *Xotodon foricurvatus*, *Promacrauchenia antiqua*, and *Cyonasua argentina?*
 13. We regard here the first records shared with the Chasicoan as lineages that originated during this time and perdurated in younger times. Quite the contrary, the numerous first records shared with Huayquerian localities establish a minimum age for the «Conglomerado osífero» (which is here considered older than the type Montehermosan with which shares two taxa; see above). We find particularly significant the first record in South America of the family of Holarctic origin Procyonidae, which is perhaps represented by the same species both in Catamarca and Entre Ríos (*Cyonasua argentina*; Marshall and Patterson, 1981). *Cyonasua* is the first immigrant from North America of the major biogeographic late Cenozoic event named "Great American Biotic Interchange." This event deeply transformed the South American mammal fauna: presently about 50% of species is Holarctic in origin. Besides, some other taxa of higher than genus level appear for the first time in the Huayquerian and the "Conglomerado osífero:" Myocastoridae, Abrocomidae, Neuryurini, Doedicurinae.
 14. Consequently, the significative secondary mixture of Pansantacrucian, Araucanian and Panpampean taxa (see Reig, 1957; Pascual and Odreman Rivas, 1971; Marshall *et al.*, 1983) present in the «Conglomerado osífero» is not sustained. We regard that the «Conglomerado osífero» (and the fauna enclosed) does not represent a long lapse. With the present evidence, the «Conglomerado osífero» appears to be Huayquerian in age.
 15. The base of Huayquerian is located at about 9 Ma (Flynn and Swisher, 1995). According to Flynn and Swisher (1995), the available radioisotopic dates and location of the boundary within the early part of Chron C3Ar, and using chron terminology and the time scale of Cande and Kent (1995) and Berggren *et al.* (1997), the best estimate for the Huayquerian/Montehermosan boundary is about 6.8 Ma, and certainly older than 6.5 Ma. Actually, this estimate is not for the boundary but to the youngest Huayquerian beds known. Beds overlying the Andalhuala Formation possibly are Chapadmalalan in age (see Cione and Tonni, 1996). Flynn and Swisher (1995) indicate that the Huayquerian to Montehermosan interval is relatively well sampled paleomagnetically in Bolivia and northwestern Argentina. However, the «Montehermosan» sections might be Chapadmalalan in age (Cione and Tonni, 1996).

16. The Huayquerian is so correlated with the Tortonian of the Geologic Time Scale (11.2 to 7.1 Ma; Gradstein and Ogg, 1996).
17. The «Conglomerado osifero» does not share *Proeuphractus* and *Vetelia* with the Chasicóan. These taxa occur in the Huayquerian beds at the Chasicó area that were considered as an early Huayquerian (*Macrochorobates scalabrinii* Biozone; Tonni et al., 1998). However, it is too premature to attribute a younger age than *Macrochorobates scalabrinii* Biozone to the «Conglomerado osifero.»
18. The presence of *Cyonasua* permits to correlate approximately the «Conglomerado osifero» and the Andalhuala Formation with those North American beds of Hemphillian age where the first South American immigrants occur for the first time (Marshall, 1985 but see Pascual et al., 1965 for a different view).

Correlation with the «Puelchan»

The «Formación» Puelche (or informally the «Puelchan») occurs deep in the subsoil of the Pampean region. The «Puelchan» is today based on thick, water-saturated, fluvial sands (and only known by drilling; Sala and Auge, 1970; Pascual and Odreman Rivas, 1971). The «Puelchan» also overlies the marine Paraná Formation (Middle Miocene-earliest Late Miocene, Cione, 1988). Santa Cruz (1972) formally designated this unit as the Puelche Formation. This name is not acceptable for it is not geographic; it refers to an Indian tribe from Patagonia. However, it is worthy of formal recognition (following North American Commission on Stratigraphic Nomenclature, 1983, p. 851) for its uniqueness and economic importance. All the fossils that are known from this unit have been obtained from drilling operations by means of suction pumps, without any kind of stratigraphic control (see synthesis of Rusconi, 1948, 1949, and also see Castellanos, 1936). In the same unit, marine vertebrates typical of the marine Paraná Formation and the «Conglomerado osifero» occur: *Carcharias acutissima*, *Isurus hastalis*, *Carcharhinus* spp., Myliobatoidei and Sciaenidae (Rusconi, 1948; personal observation ALC) along with terrestrial vertebrates (toxodontid and rodents; see Rusconi, 1948, 1949; Cione and Tonni, 1995; Bond, 1999; Bond et al., 1995; Bond and López, 1998). However, these remains formed the basis for speculating that the mammals comprised an ancient pampean fauna that could well be «Ensenadense basal» (Pascual et al., 1965) or older, but always pampean (Simpson, 1940). This is despite the fact that «mammal remains from the Puelche are, in most cases, fragmentary and their determination is according to Pascual et al. (1965, p. 180), more doubtful than believed by Rusconi» (Marshall et al., 1984, p. 24). Pascual et al. (1965) established that the fauna had a post-Chapadmalalan aspect and included Ensenadan taxa. Despite the lack of stratigraphic control, Pascual et al. (1965), Reig (1981), and Marshall et al. (1984; and all other workers) included the mammals from the «Puelchense» in the «Uquian» faunal list, and even correlated the «Puelchense» with the lower part of «Uquian Land-mammal age» (cf. Kraglievich, 1952; Reig, 1981).

We consider that the presence of the abundant Miocene vertebrates and the stratigraphic relationships confirm the correlation of at least the base of the Puelchan beds with the Ituzaingó Formation. We identify no Marplatán (the age post-Chapadmalalan and pre-Ensenadan that replaces the «Uquian age;» see Cione and Tonni, 1995) from the Puelchan beds.

Biogeography

Marine vertebrates

Genera of marine fishes in the Paraná Formation are those known from the warmer part of the Argentinian Biogeographic Province (*sensu* López, 1964; see Cione, 1978, 1988). The fauna is integrated by inhabitants of the shelf, probably inner shelf. No typical oceanic taxon was found yet.

Remarkable is the presence of heterodontid sharks, which presently do not inhabit the Atlantic Ocean. Other sharks such as pristiophorids were detected in late Oligocene-Middle Miocene beds in Patagonia. *Heterodontus* and pristiophorid species were almost worldwide in the Tertiary (Cappetta, 1987; Cione, 1988; Cione and Expósito, 1980). *Heterodontus* species live in the Pacific today (Compagno, 1984), whereas pristiophorids do not inhabit modern South American Atlantic and Pacific coasts (Compagno, 1984). Presently, only a small population of the endemic species *Pristiophorus schroederi* occurs in the (northwestern) Atlantic (Springer and Bullis, 1960). Both groups greatly reduced their distribution after the Miocene. The finding of *Pristiophorus* in the Puerta del Diablo Formation of Patagonia and *Heterodontus* in the Paraná Formation constitute the last records of these genera in the South Atlantic and in the Atlantic, respectively. Both represent an example of extirpation in a particular area and persistence in other (Cione and Azpelicueta, in preparation).

The cetaceans Pontoporiidae were also common in the northern South Pacific (north of Chile and south of Perú, Cozzuol, 1996), in the North Pacific (Barnes, 1985) and North Atlantic. The presence of pontoporiids in the Paraná Formation suggest a biogeographical connection with those regions, in a time in which the Central American seaway was still open.

Dioplotherium, apparently present in those deposits, was also found in northern Brazil (Pirabas Formation, de Toledo and Domning, 1991) and in the North Atlantic and Pacific (Domning, 1989). Balaenopterids were proposed to be originated in the tropical Atlantic not before than late Middle Miocene, but they appeared worldwide in the record by the Late Miocene, as is the case of Paraná Formation.

If the *Monachus* affinities of *Poperiptychus argentinus* are confirmed, this group is also known to have a tropical distribution.

Consequently, aquatic mammals indicates stronger connections South-North than with others localities of the southern Hemisphere.

Aquatic continental vertebrates

The Neotropical Region is divided into two phenetic subunits according to recent fish distribution (Ringuélet, 1975; Arratia *et al.*, 1983): the Brazilian and Austral subregions. The Austral Subregion includes Patagonia, Cuyo, and southern-central Chile (Ringuélet, 1975; Arratia *et al.*, 1983; Arratia, 1997). The northern boundary of the Austral subregion is defined in Argentina by the San Juan-Desaguadero-Curacó-Colorado drainages. The Brazilian Subregion occupies the rest of South America and Central America. The Brazilian Subregion is the richest area in fish diversity and the Austral Subregion is one of the less diverse areas in the world (less than 25 species; Ringuélet, 1975; Almirón *et al.*, 1997; Arratia, 1997; Casciotta *et al.*, 1999). The Brazilian diversity dramatically diminishes south of the Río de la Plata today.

Miocene freshwater fishes from the Paraná area are typically brazilic and no austral taxon was detected.

Fishes, crocodiles, cetacean, and sirenids from the “Conglomerado osífero” are related to those of the late Miocene Laventan beds (Langston, 1965; Langston and Gasparini, 1997; Cione,

Lundberg and Machado Allison, en preparation), the Late Miocene beds of the Peru and Brazilian Acre (Boquentin *et al.*, 1989; Rancy *et al.*, 1989; Bocquentin and Souza Filho, 1990; Frailey, 1986; Gasparini, 1996) and the late Miocene–Lower Pliocene beds of Urumaco, Venezuela (Buffetaut, 1982; Gasparini, 1996). La Venta, Acre and Paraná share the tetrapods *Caiman*, *Mourasuchus* and *Gryposuchus*. Acre and Paraná also share the tetrapods *Ischyrorhynchus*, *Sauroctes* and *Ribodon*.

Presently, the Paraná-Uruguay-Plata basin is practically isolated from the Amazonian basin. The presence of similar freshwater aquatic vertebrates with Miocene beds from Colombia (present Magdalena basin), Venezuela and the Brazilian Amazonia seems to indicate hydrographic relationships that does not exist today. The occurrence of iniiid cetaceans, trichechid sirenians and a very high diversity of crocodiles also is suggestive of warmer climate than present.

Terrestrial vertebrates

Terrestrial vertebrates show a very high diversity both taxic and in inferred habits. The heterogeneous landscape, controlled by one or several rivers favoring the development of varied vegetation in a reduced area and the presence of neighbouring open areas could explain that diversity.

Temperatures and humidity were higher in the whole central and northwestern Argentina area what is clearly evidenced by the fauna and flora found from the “Araucanense,” the Huayquerian beds of the Pampean and Mesopotamic areas. However, among the terrestrial vertebrates, xenarthran and notoungulate taxa indicate that the Mesopotamic area was already a biogeographic area different to that of the Pampean and northwestern Argentina (Scillato Yané, 1975b), more closely related to Uruguay and the Brazilian Acre areas (see below) at least since the Late Miocene. Consequently, the Guayano-Brazilic dominion, as it is known today, it was probably beginning to develop. However, rodents, xenarthrans, ungulates, marsupials and carnivores showed also affinities with Catamarcan localities. The typical Central or Subandean Dominion of the Andean-Patagonian Subregion (Ringuelet, 1961; Figure 9) was not yet beginning to differentiate.

The absence of some taxa which are recorded in coeval sediments of relatively nearby regions (Glyptodontinae glyptodonts, and Myrmecophagidae and Cyclopidae vermilinguans) is noteworthy. Notoungulates and caviomorphs showed also high endemism. During the Miocene the zoogeographic and environmental characteristics of this part of the Mesopotamia must have been quite peculiar. This hypothesis is not hindered by the analysis of the rest of the biota and the geological information. With the available evidence, the closest affinities of the «Mesopotamian» xenarthrans may be found within those of the Miocene-Pliocene of Uruguay.

The “Conglomerado osífero” fauna is also important for studying the evolution of birds in relation to the “Great American Biotic Interchange” because it corresponds to a older moment than the establishment of a definitive connection between South and North America. The fossil record shows that flightless birds (and those with reduced capacity of flight) such as Rheiformes, Tinamiformes, “phororacoids,” and Opisthocomidae evolved in complete isolation during most of the Tertiary (Tambussi and Noriega, 1996). Besides this, some flying birds such as Teratornithidae and the “suboscines” passeriforms did not cross the Panama gap until the isthmus was established (Tambussi and Noriega, 1996; Noriega, 1998). Other birds (eg dendrochenin anatids and paleodin flamingos) were not biogeographically isolated. This late evidence confirms the hypothesis favoring a significative biogeographic connexion of South American bird faunas with those of North America and Europe (Martin, 1983; Rasmussen and Kay, 1992).

Environment

Marine

The high marine level of the middle Miocene sea permitted the ingression of marine waters in the Chacopampean plains at least to Paraguay and Bolivia (Uliana and Biddle, 1988; Marshall *et al.*, 1993; Cione and Cozzuol, in preparation). In the south, the marine influence was restricted to northeastern Patagonia. A warm temperate Miocene ichthyofauna occurs in the top of the Paraná Formation. The assemblage from Paraná is different from the Patagonian ichthyofauna but it is similar to that present at the same latitude in the Atlantic coast of southern Brazil (Cione, 1978). Whereas at Paraná carcharhinids, hemigaleids, and odontaspidids dominate, Patagonian ichthyofaunas are ruled by lamnids. Cetacean from the Paraná Formation also do not indicate temperatures very different to that present in the same latitude. However, the sirenian, the pinniped and the invertebrates suggest warmer waters for Paraná, which can be extended to northern Patagonia on the basis of invertebrates (Del Río, 1988).

Remarkably, contrasting with the late Oligocene-early Miocene ichthyofaunas, Middle Miocene shark assemblages appears to have been poorly diversified in Patagonia (Cione, 1978, 1988; Cione and Tonni, 1981; Perea *et al.*, 1985; Perea and Ubilla, 1989, 1990).

The elasmobranch association present in the Paraná Formation suggest normal marine salinity.

Continental

Neither dipnoan nor anuran were recorded in the area. Taking into consideration the relatively good preservation of dipnoan teeth, lack of these fishes could be related to the absence of appropriate lentic environments (but see below evidence from birds). The absence of anuran could be due to a defect of preservation and collection.

The paleogeographic location of Paraná, in the subtropical to temperate belt is also confirmed by the absence of crocodylids, podocnemidids, and primates. Today, South American crocodylids inhabit strictly tropical regions. However, the lack of primates would be due to insufficient preservation.

The taxonomic reptile diversity, especially crocodiles from the «Conglomerado osífero» suggest varied paleoenvironments. The predominance of aquatic birds in the «Conglomerado osífero» support the presence of woody lowlands and swamps along the river banks. This type of environments is also a requisite for the trichechids and iniids, which need lakes associated to the main river to live, feed and reproduce. Besides, rheas and fororhacs suggest a savanna-like environments near the riversides (Noriega, 1994, 1995).

Glyptodontids are very diversified in the «Conglomerado osífero.» The glyptodontid diversity from the «Araucanian» of Catamarca and Tucumán is smaller. However, the main difference between both faunas lies in the very scarce glyptodontid frequency in Entre Ríos in comparison with their extraordinary abundance in the Northwest of Argentina. As middle to large sized mammals, all these glyptodonts must have preferred more open environments (savannas, grasslands, and herbaceous steppes). These conditions were probably common in the northwest but not in the Paraná area. Here, glyptodonts are represented mainly by isolated scutes, small fragments of carapace or skeleton, with clear evidence of post-mortem transportation. This suggests that they lived near the gallery forests of the «pre-Paraná», but not into them.

The tardigrade diversity must have been related to a heterogeneous landscape, surely

controlled by a river (or several rivers), favoring the development of a varied vegetation cover in reduced areas.

Comparing with other faunas from the Late Miocene and Pliocene of South America, protheriids are extremely abundant in the "Conglomerado osífero," confirming the relation of these mammals to woody and wet environments. Interatheriidae Interatheriinae notoungulates also seem to have been related to woody and wet environments while Hegetotheriidae, especially the "rodentiform" Pachyrukhinae would have inhabited more open and dry environments. An analogous situation was discovered in the Colloncuran faunas of Argentina and Laventan faunas of Colombia where Pachyrukhinae were very abundant in Patagonia, although Interatheriinae were also present. On the contrary, in Colombia, where the environment seem to have been more tropical, woody and wet, Pachyrukhinae are absent (Cerdeño and Bond, 1998) and only an endemic and large taxon of Interatheriinae, though smaller than *Munizia* occur.

The occurrence of a porcupine rodent support the presence of forested environments, as well as the biogeographic connexion with northern South America. The absence of octodontid rodents suggest the possible absence of arid environments. Octodontids are common in western and central Argentina (Central or Subandean Dominion).

Procyonids are medium-sized, good swimmers omnivores that mainly inhabit forested environments (Bond, 1986).

Among the terrestrial vertebrates, no primate was recorded. The lack of monkeys would be due to a defect of preservation because global temperatures were higher than present, primates had been recorded until Colloncuran times in Patagonia, forested areas certainly occurred in the area of Paraná (evidenced by the abundant wood present in the sediments and the association of terrestrial vertebrates) and other arboreal mammals such as porcupines are now known in the "Conglomerado osífero."

Acknowledgments

The authors would like to thank the following institutions and people:

Consejo Nacional de Investigaciones Científicas y Tecnológicas, Agencia Nacional de Promoción Científica y Tecnológica, Comisión de Investigaciones Científicas de la Provincia de Buenos Aires, Facultad de Ciencias Naturales y Museo for permanent financial support.

Centro de Investigaciones Científicas y Transferencia de Diamante, for assistance in the area.

F. Aceñolaza and R. Herbst, for the invitation to participate in the present volume. C. Ceruti, for assistance in Paraná. José Laza, for preparation of materials. Cecilia Deschamps for assistant to some of the authors. Carlos Tremouilles, for drawings. Carlos Steger, for loaning valuable materials.

Subfamily Protheroetheriinae									
<i>Protheroetherium</i> Ameghino, 1883								X	
<i>P. cervioides</i> Ameghino, 1883								X	
<i>P. gradatum</i> Ameghino, 1891								X	
<i>Epitherium</i> Ameghino, 1888								X	
<i>E. paranensis</i> (Ameghino, 1904)								X	
<i>Brachytherium</i> Ameghino, 1883					?		X	X	
<i>B. cuspidatum</i> Ameghino, 1883								X	
<i>Licaphrium</i> Ameghino, 1887								X	
<i>L. mesopotamiense</i> Bianchini y Bianchini, 1971								X	
<i>Thoatherium</i> Ameghino, 1887								?	
? <i>Thoatherium</i> sp.								?	
<i>Coelosoma</i> Ameghino, 1891								X	
<i>C. eversa</i> Ameghino, 1891								X	
Family Macraucheniiidae									
Subfamily Macraucheniiinae									
<i>Scalabrinitherium</i> Ameghino, 1883								X	
<i>S. bravardi</i> Ameghino, 1883								X	
<i>S. rothi</i> Ameghino, 1885								X	
<i>Oxydontherium</i> Ameghino, 1883								X	
<i>O. zeballosi</i> Ameghino, 1883								X	
<i>Mesorhinus</i> Ameghino, 1885								X	
<i>M. piramydatus</i> Ameghino, 1885								X	
<i>Paranauchenia</i> Ameghino, 1904								X	
<i>P. denticulata</i> (Ameghino, 1891)								X	
<i>Promacrauchenia</i> Ameghino, 1904							X	X	HMC
<i>P. antiqua</i> Ameghino, 1887								X	M
<i>Cullinia</i> Cabrera y Kraglievich, 1931				X				?	
<i>Cullinia</i> sp.								?	

TABLE 1. Stratigraphic and geographic range of xenarthran taxa. Abbreviations: CC, Colloncuran; May, Mayoan; Lav, Laventan; ChV, El Vivero Member of the Arroyo Chasicó Formation; ChB, Las Barrancas Member of the Arroyo Chasicó Formation; Itz, "Conglomerado osífero" of the Ituzzaingó Formation; CTAr, Andalhualá Formation; Ur, Miocene-Pliocene beds of Uruguay; H, Epecuén "Formation" and Cerro Azul Formation of Pampean area; M, Montehermosan localities of Pampean area; C, Chapadmalalan localities of Pampean area.

Caviomorph Rodents	CC	Lav	Ch. V	Ch. B	Itz	CT Ar	MP Ur	HMC
Family Dinomyidae								
<i>Briaromys</i> Ameghino, 1889					X			
<i>Carlesia</i> Kraglievich, 1926			X		X	X		
<i>Diaphoromys</i> Kraglievich, 1931			?	?	X			H
<i>Doellomys</i> Alvarez, 1947					X			
<i>Eumegamys</i> Kraglievich, 1926					X			
<i>Eumegamysops</i> Alvarez, 1947					X			
<i>Gyriabrus</i> Ameghino, 1891			?	?	X		X	H?
<i>Isostylomys</i> Kraglievich, 1926					X		X	
<i>Paranamys</i> Kraglievich, 1934					X			
<i>Pentastylodon</i> Alvarez, 1947					X			
<i>Pentastylomys</i> Kraglievich, 1926					X			
<i>Potamarchus</i> Burmeister, 1885			?	?	X			H?
<i>Pseudosignomys</i> Kraglievich, 1931					X			
<i>Protomegamys</i> Kraglievich, 1932					X			
<i>Tetrastylomys</i> Kraglievich, 1926					X			H
<i>Tetrastylus</i> Ameghino, 1886			X		X	X		H
Family Chinchillidae								
<i>Lagostomopsis</i> Kraglievich, 1926			?	X	X	X	X	HMC
Family Neoepilemidae								
<i>Phoberomys</i> Kraglievich, 1926					X			
<i>Perimys</i> Ameghino, 1887					?			
<i>Neoepilema</i> Ameghino, 1889					X			
Family Abrocomidae								
<i>Protabrocoma</i> Kraglievich, 1927					X			H
Family Echimyidae								
<i>"Eumysops" parodii</i> Kraglievich, 1945					X			
<i>Haplostropha</i> Ameghino, 1891					X			
Family Erethizontidae								
<i>Paradoxomys</i> Ameghino, 1886					X			
Family Caviidae								
<i>Cardiomys</i> Ameghino, 1885	?			X	X	X		HMC
<i>Caviodon</i> Ameghino, 1885					X			HMC
<i>Palaococavia</i> Ameghino, 1889					X	X		HMC
<i>Parodimys</i> Kraglievich, 1932			X		X			
<i>Pliodolichotis</i> Kraglievich, 1927					X			
<i>Prodolichotis</i> Kraglievich, 1932		X			X	X		
Family Hydrochoeridae								
<i>Anatochoerus</i> Mones & Vucetich, 1991					X			
<i>Anchimys</i> Ameghino, 1886					X			
<i>Anchimysops</i> Kraglievich, 1927					X			M
<i>Cardiatherium</i> Ameghino, 1883			?	?	X			H
<i>Contraocavia</i> Burmeister, 1885					X			
<i>Kiyutherium</i> Francis & Mones, 1965					X	X	X	H
<i>Plexochoerus</i> Ameghino, 1886					X			
<i>Procardiatherium</i> Ameghino, 1885				X	X			H
<i>Protohydrochoerus</i> Rovereto, 1914					X			MC
Family Myocastoridae								
<i>Colpostemma</i> Ameghino, 1891					X			
<i>Myocastor</i> Kerr, 1792					?			

TABLE 2. Stratigraphic and geographic range of "native ungulate" taxa.

Xenarthrans	CC	May	Lav	Ch. V	Ch. B	Itz	CT Ar	MP Ur	HMC
Order Cingulata Superfamily Dasypodoidea Family Dasypodidae Subfamily Dasypodinae Tribe Dasypodini <i>Dasypus</i> Linneé, 1758 <i>D. neogaeus</i> (Ameghino, 1891)						X X			
Subfamily Euphractinae Tribe Eutatini <i>Chasicotatus</i> Scillato-Yané, 1977 <i>Ch.</i> sp. nov.					X	X X	X		
Tribe Euphractini <i>Macroephractus</i> Ameghino, 1887 <i>M. retusus</i> Ameghino, 1887 <i>Proephractus</i> Ameghino, 1886 <i>P. limpidus</i> Ameghino, 1886 <i>Zaedyus?</i> Ameghino, 1889 <i>Z.</i> sp.					X	X X X X X	X	X X	HMC MC
Superfamily Glyptodontoidea Family Pampatheriidae <i>Kraglievichia</i> Castellanos, 1927 <i>K. paranense</i> (Ameghino, 1883) <i>Scirrotherium</i> Edmund & Theodore, 1997 <i>S.</i> sp. nov.			X		X	X X X X		X X	
Family Glyptodontidae Subfamily Sclerocalyptinae Tribe Sclerocalyptini <i>Berthawileria</i> Castellanos 1939 <i>B.</i> sp. Sclerocalyptini inc. "Hoplophorus" Ameghino 1889, non Lund « <i>H.</i> » <i>verus</i> Ameghino, 1889						X X X		X	C
Tribe Lomaphorini <i>Trachycalyptus?</i> Ameghino, 1908 <i>T?</i> <i>cingulatus</i> (Ameghino, 1889)						X			
Tribe Palaehoplophorini <i>Palaehoplophorus</i> Ameghino, 1883 <i>P. antiquus</i> (Ameghino, 1883) <i>Chlamyphractus</i> Castellanos, 1939 <i>Ch. pressulus</i> (Ameghino, 1885) <i>Protoglyptodon</i> Ameghino, 1885 <i>P. primiformis</i> Ameghino, 1885	X					X X X X X X			
Tribe Plohophorini <i>Plohophorus</i> Ameghino, 1887 <i>P. paranensis</i> Ameghino, 1891 <i>Parahoplophorus</i> Castellanos, 1931-32 <i>P. paranensis</i> (Ameghino, 1883)						X X X X	X	X	H? MC
Tribe Neuryurini <i>Urotherium</i> Castellanos, 1926 <i>U. interundatum</i> (Ameghino, 1885)						X X	X	X X	C

Bibliography

- Aceñolaza, F.G. 1976. Consideraciones bioestratigráficas sobre el Terciario marino de Paraná y alrededores. *Acta Geológica Lilloana* 13: 91-107.
- Aceñolaza, P.G. and Aceñolaza, F.G. 1996. Improntas foliares de una Lauraceae en la Formación Paraná (Mioceno superior), en la Villa Urquiza, Entre Ríos. *Ameghiniana* 33: 155-159.
- Aceñolaza, F.G. and Aceñolaza, G. 1999. Trazas fósiles del Terciario marino de Entre Ríos (Formación Paraná, Mioceno medio), República Argentina. *Boletín de la Academia Nacional de Ciencias*, Córdoba 64: 209-233.
- Aceñolaza, F.G. and Sayago, J.M.. 1980. Análisis preliminar sobre la estratigrafía, morfodinámica y morfogénesis de la región de Villa Urquiza, provincia de Entre Ríos. *Acta Geológica Lilloana* 15: 139-154.
- Albino, A. 1996. The South American fossil Squamata (Reptilia: Lepidosauria). In: G. Arratia (ed.), *Contributions of Southern South America to Vertebrate Paleontology*, München Geowissenschaft Abhandlungen 30, pp. 185-202.
- Alessandri, G. 1896. Ricerche sui pesci fossili de Paraná. *Atti della Reale Accademia di Scienze di Torino* 31: 1-17.
- Almirón, A., Azpelicueta, M.M., Casciotta, J. and López Cazorla, A. 1997. Ichthyogeographic boundary between the Brazilian and Austral subregions in South America, Argentina. *Biogeographica* 73: 23-30.
- Ambrosetti, J. 1890. Observaciones sobre los reptiles fósiles oligocenos de los terrenos terciarios antiguos del Paraná. *Boletín Academia Nacional de Ciencias de Córdoba* 10: 409-426.
- Ambrosetti, J. 1893. Contribución al estudio de las tortugas oligocenas de los terrenos terciarios antiguos del Paraná. *Boletín Instituto Geográfico Argentino* 14: 489-499.
- Ameghino, F. 1883a. Sobre una colección de mamíferos fósiles del Piso Mesopotámico de la Formación Patagónica recogidos en las barrancas del Paraná por el Profesor Pedro Scalabrini. *Boletín de la Academia Nacional de Ciencias de Córdoba* (República Argentina) 5 (entrega 1a): 101-116.
- Ameghino, F. 1883b. Sobre una nueva colección de mamíferos fósiles recogidos por el Profesor Scalabrini en las barrancas del Paraná. *Boletín de la Academia Nacional de Ciencias de Córdoba* 5: 112-113.
- Ameghino, F. 1885. Nuevos restos de mamíferos fósiles oligocenos recogidos por el profesor Pedro Scalabrini y pertenecientes al Museo Provincial de la ciudad de Paraná. *Boletín de la Academia Nacional de Ciencias de Córdoba* 8: 3-207.
- Ameghino, F. 1886. Contribución al conocimiento de los mamíferos fósiles de los terrenos Terciarios antiguos del Paraná. *Boletín de la Academia Nacional de Ciencias de Córdoba* (República Argentina) 9: 3-226.
- Ameghino, F. 1887. Observaciones generales sobre el orden de los mamíferos extinguidos llamados toxodontes (Toxodontia) y sinopsis de los géneros y especies hasta ahora conocidos. *Anales del Museo de La Plata*, 1 (entrega 1): 1-66.
- Ameghino, F. 1889. Contribución al conocimiento de los mamíferos fósiles de la República Argentina. *Actas de la Academia Nacional de Ciencias de Córdoba*. 6: XXXII + 1028, Atlas: 98 láms.
- Ameghino, F. 1891. Caracteres diagnósticos de cincuenta especies nuevas de mamíferos fósiles argentinos. *Revista Argentina de Historia Natural* 1: 129-167.
- Ameghino, F. 1892. Répliques aux critiques du Dr. Burmeister sur quelques genres de mammifères fossiles de la République Argentine. *Boletín de la Academia Nacional de Ciencias de Córdoba* (República Argentina), 12 (Entrega 4a): 437-469.
- Ameghino, F. 1898. Sinopsis geológico-paleontológica. En: *2do Censo de la Republica Argentina. I. Territorio*, Buenos Aires: 1-500.
- Ameghino, F. 1904a. Recherches de morphologie phylogénétique sur les molaires supérieures des ongulés. *Anales del Museo Nacional de Buenos Aires* 9 (ser. 3a, 3): 1-541.
- Ameghino, F. 1904b. Nuevas especies de mamíferos, cretáceos y terciarios de la República Argentina. *Anales de la Sociedad Científica Argentina* 20: 56-58.
- Ameghino, F. 1906. Les Formations Sédimentaires du Crétacé Supérieur et du Tertiaire de Patagonie avec un parallèle entre leurs faunes mammalogiques et celles de l'ancien continent. *Anales del Museo Nacional de Buenos Aires* 15 (ser. 3a, 8): 1-568.
- Arratia, G., Peñafort, B. and Menu-Marque, S. 1983. Peces de la Región Sureste de los Andes y sus probables relaciones biogeográficas actuales. *Deserta* 7: 48-107.
- Arratia, G. and Cione, A.L. 1996. The fossil fish record of Southern South America. In: G. Arratia (ed.), *Contributions of Southern South America to Vertebrate Paleontology*, München Geowissenschaft Abhandlungen 30, pp. 9-72.
- Arratia, G. 1997. Brazilian and Austral freshwater fish faunas of South America. A contrast. In: *Tropical biodiversity and systematics*. Proceedings of the International Symposium on Biodiversity and Systematics in Tropical Ecosystems, Bonn, 1994: 179-186.
- Báez, A.M. and Gasparini, Z. 1977. Orígenes y evolución de los anfibios y reptiles del Cenozoico de América del Sur. *Acta Geológica Lilloana* 14: 149-232.
- Berggren, W.A., Hilgen, F.J., Langereis, C.G., Kent, D., Obradovich, J., Raffi, I., Raymo, M.E., and Shackleton, N.J. 1997. Late Neogene chronology: new perspectives in high-resolution stratigraphy. *Bulletin of the Geological Society of America* 107: 1272-1287.
- Bianchini, L.H. Delupi de, and Bianchini, J.J. 1971. Revisión de los Protheroheriinae (Mammalia, Litopterna) del Mesopotamiense. *Ameghiniana* 8: 1-24.

- Bocquentin, J. and Souza Filho, J. 1990. O crocodiliano sud-americano *Carandaisuchus* como sinonimia de *Mourasuchus* (Nettosuchidae). *Revista Brasileira de Geociências* 20: 230-233.
- Bond, M. 1986. Los ungulados fósiles de Argentina. *Actas del IV Congreso Argentino de Paleontología y Biostratigrafía*. Mendoza: 173.
- Bond, M. 1999. Quaternary native ungulates of Southern South America. A synthesis. *In: E. P. Tonni and A. L. Cione (eds.). Quaternary vertebrate paleontology in South America*. Quaternary of South America 12. Special Volume, pp. 177-205.
- Bond, M. and López, G.M. 1996. Los Macraucheniiidae (Mammalia, Litopterna) de la Formación Arroyo Chasicó (Partido de Villarino, Pcia. de Buenos Aires). *Actas Cuartas Jornadas Geológicas y Geofísicas Bonaerenses*, Junín 1: 23-27.
- Bond, M. and López, G.M. 1998. *Notopachyrukhos tambutoi* Rusconi, 1933, un Intertheriidae (Mammalia, Notoungulata) del Puelchense de la provincia de Buenos Aires. *Actas del X Congreso Latinoamericano de Geología y VI Congreso Nacional de Geología Económica Buenos Aires* 1: 280-283.
- Bond, M.; Cerdeño, M.E. and López, G.M. 1995. Ungulados Nativos Sudamericanos. En: M. T Alberdi, G. Leone and E. P. Tonni (eds.). *Evolución Biológica y Climática de la región Pampeana durante los últimos cinco millones de años. Un ensayo de correlación con el Mediterráneo Occidental*. Monografías del Museo Nacional de Ciencias Naturales, Madrid, CSIC, pp. 257-275.
- Bondesio, P., Laza, J., Scillato Yané, G., Tonni, E.P. and Vucetich, M.G. 1980. Estado actual del conocimiento de los vertebrados de la Formación Arroyo Chasicó (Plioceno temprano) de la provincia de Buenos Aires. *Actas II Congreso Argentino de Paleontología y Biostratigrafía y I Congreso Latinoamericano de Paleontología* 3: 101-127.
- Bravard, A. 1858. Monografía de los terrenos terciarios del Paraná. (Reimpresión facsimilar con prólogo de F.G. Aceñolaza, Imprenta del Congreso de la Nación, Buenos Aires, 1995, V-XII + 1-107 pp.).
- Broin, F. de and de la Fuente, M. 1993. Les tortues fossiles d'Argentine: Synthèse. *Annales de Paléontologie* 79: 169-231.
- Buffetaut, E. 1982. Systématique, origine et évolution des Gavialidae Sud-Américains. *Geobios* 6: 127-140.
- Burmeister, G. 1871. On *Saurocetes argentinus*, a new type of Zeuglodontidae. *Annals Magazine of Natural History*, London 47: 51-55.
- Burmeister, G. 1885. Examen crítico de los mamíferos y los reptiles denominados por Don Augusto Bravard. *Anales del Museo Público de Buenos Aires* 3: 95-173.
- Burmeister, G. 1891. IX. Continuación a las adiciones al examen crítico de los mamíferos fósiles terciarios para dar a conocer los objetos de historia natural nuevos ó poco conocidos conservados en ese establecimiento. *Anales del Museo Nacional de Buenos Aires* 3: 401-461.
- Butler, R.F., Marshall, L.G., Drake, R.E. and Curtiss, G.H. 1984. Magnetic polarity stratigraphy and 40K-40Ar dating of Late Miocene and early Pliocene continental deposits, Catamarca Province, NW Argentina. *Journal of Geology* 92: 623-636.
- Cande, S.C. and Kent, D. 1995. Revised calibration of the Geomagnetic Polarity Time Scale for the Late Cretaceous and Cenozoic. *Journal of Geophysical Research* 100: 6093-6096.
- Capetta, H. 1987. Chondrichthyes II. In: H.-P. Schultze (ed.), *Handbook of Paleichthyology*. Gustav Fischer Verlag, Stuttgart, New York, pp. 1-193
- Carlini A.A., Scillato-Yané G.J. and Vizcaíno, S.F. 1990. Un singular Paratheria del Eoceno temprano de Patagonia, Argentina. *VII Jornadas Argentinas de Paleontología de Vertebrados*, Buenos Aires, 1990. *Ameghiniana* 26: 241.
- Carlini, A.A., Ortiz-Jaureguizar, E., Pascual, R., Scillato-Yané, G.J. and Vizcaíno, S.F. 1994. The negative paleontological record on the controverted origin and relationships of the Xenarthra. *Resúmenes del VI Congreso Argentino de Paleontología y Biostratigrafía* Trelew 21-22.
- Carlini, A.A., Pascual, R., Reguero, M., Scillato-Yané, G.J., Tonni, E.P. and Vizcaíno, S.F. 1990. The first paleogene land placental mammal from Antarctica: its paleoclimatic and paleobiogeographical bearings. *IV International Congress of Systematic and Evolutionary Biology*. Washington. *Evolutionary Biology* 1990
- Carlini A.A, Heizler, M.H., Kay, R.F, Madden, R.H., Mazzoni, M.M., Re, G.H., Sandeman, H. and Vucetich, M.G. 1999. Implications of the Late Eocene Age of the Casamayoran fauna at Gran Barranca, Chubut, Argentina. *Journal of Vertebrate Paleontology* 19, Supplement 3: 35A.
- Carlini, A. A, Scillato- Yané, G. J., Noriega, J. I. y Aceñolaza, F. In press. Perezosos terrestres (Xenarthra, Tardigrada) del «Mesopotamiense» (Formación ituzzaingó, Mioceno tardío-Plioceno temprano) de la Provincia de Entre Ríos, Argentina. *Sivdia Geológica Salmanticensis*, 36: 13-27. Salamanca.
- Casciotta, J., Almirón, A., Cione, A. y Azpelicueta, M. M. 1999. Brazilian freshwater fish assemblages from southern Pampean area, Argentina. *Biogeographica* 75: 67-78.
- Castellanos, A. 1936. Los sedimentos del Pampeano inferior y del Araucanense en el valle de los Reartes (Sierra de Córdoba). *Publicaciones Facultad de Ciencias Matemáticas, Físicoquímicas y Naturales* 6: 1-16.
- Cerdeño, E. and Bond, M. 1998. Taxonomic revision and phylogeny of *Paedotherium* and *Tremacyllus* (Pachyrukhinae, Hegetotheriidae, Notoungulata) from the Late Miocene to the Pleistocene of Argentina. *Journal of Vertebrate Paleontology* 18: 799-811.
- Chebli, G.A., Tofalo, O. and Turzzini, G. 1989. Mesopotamia. *In: G.A. Chebli and L. Spalletti (eds.), Cuencas sedimentarias argentinas*. San Miguel de Tucumán, Universidad Nacional de Tucumán, Instituto de Correlación Geológica, Serie Correlación Geológica 6, pp. 79-100.
- Cione, A.L. 1978 Aportes paleoictiológicos al conocimiento de la evolución de las paleotemperaturas en el área austral del América del Sur durante el Cenozoico. Aspectos zoogeográficos y ecológicos conexos. *Ameghiniana* 15: 183-208.

- Cione, A.L. 1986. Los peces continentales del Cenozoico de Argentina. Su significación paleoambiental y paleobiogeográfica. *Actas del III Congreso Argentino de Paleontología y Biostratigrafía* 2: 101-106.
- Cione, A.L. 1988. *Los peces de las formaciones marinas del Cenozoico de Argentina*. Tesis doctoral, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, pp. 588.
- Cione, A.L. and Expósito, S. 1980. Chondrichthyes del "Patagoniano" s.l. de Astra, Golfo de San Jorge, provincia de Chubut, Argentina. Su significado paleoclimático y paleobiogeográfico. *Congreso Argentino de Paleontología y Biostratigrafía, No. 2 and Congreso Latinoamericano de Paleontología* Buenos Aires 1, *Actas* 2: 275-290.
- Cione, A.L. and Tonni, E. 1981. Un pingüino (Aves, Spheniscidae) de la Formación Puerto Madryn (Mioceno tardío) de Chubut, Argentina. Comentarios acerca del origen, la paleoecología y zoogeografía de los Spheniscidae. *Congreso Latinoamericano de Paleontología* 2, Porto Alegre *Anais* 2: 591-604.
- Cione, A.L. and Torno, A.. 1984. Descripción y comparación de peculiares estructuras en la aleta dorsal de *Pogonias cromis* (Perciformes, Sciaenidae) y de una especie de la misma familia del Terciario tardío de Entre Ríos, Argentina. *Resúmenes VI Jornadas Argentinas de Zoología*: 178.
- Cione, A.L. and Casciotta, J.R. 1995. Freshwater teleostean fishes from the Miocene of the Quebrada de la Yesera, Salta, Northwestern Argentina. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* 196: 377-394.
- Cione, A.L. and Tonni, E.P. 1995. Chronostratigraphy and "Land-mammal ages" in the Cenozoic of southern South America: principles, practices, and the "Uquian" problem. *Journal of Paleontology* 69: 135-159.
- Cione, A.L. and Tonni, E.P. 1996. Reassessment of the Pliocene-Pleistocene continental time scale of southern South America. Correlation of the Chapadmalal with Bolivian sections. *Journal of South American Earth Sciences* 9: 221-236.
- Cione, A.L. and Casciotta, J.R. 1997. Miocene cynodontids (Osteichthyes: Characiformes) from Paraná, central eastern Argentina. *Journal of Vertebrate Paleontology* 17: 616-619.
- Cione, A.L., Vergani, G., Starck, D., and Herbst, R. 1995. Los peces del Mioceno de la Quebrada de la Yesera, provincia de Salta, Argentina. Su valor como indicadores ambientales. *Ameghiniana* 32: 129-140.
- Compagno, L. 1984. *Sharks of the World*. FAO Species Catalogue. 325 pp., Rome.
- Cordini, R. 1949. Contribución al conocimiento de la geología económica de Entre Ríos. *Anales de la Dirección General de Industrias y Minería* 2: 1-45.
- Cozzuol, M.A. 1985. The odontoceti of the "Mesopotamiense" of the Paraná River Ravines. Systematic Review. In: G. Pilleri (ed.), *Investigations on Cetacea* 17, pp. 39-54.
- Cozzuol, M.A. 1989. Una nueva especie de *Saurodelphis* Burmeister, 1891 (Cetacea: Iniidae) del "Mesopotamiense" (Mioceno tardío-Plioceno temprano) de la provincia de Entre Ríos, Argentina. *Ameghiniana* 25: 39-45.
- Cozzuol, M.A. 1993. *Mamíferos acuáticos del Mioceno medio y tardío de Argentina. Sistemática, evolución y biogeografía*. Tesis doctoral, Universidad Nacional de La Plata, Facultad de Ciencias Naturales y Museo de La Plata, inédita.
- Cozzuol, M.A. 1996. The record of the aquatic mammals in southern South America. In: G. Arratia, (ed.). *Contributions to southern South America to Vertebrate Paleontology*, Münchner Geowissenschaftliche Abhandlungen, Reihe A, Geologie und Paläontologie 30: 321-342.
- Darwin, C. 1839. *Voyage of the Beagle. Journal of researches into the Geology and Natural History of the various Countries visited by H.M.S Beagle*. Prometheus Books, London, 547 p.
- de la Fuente, M. 1988. *Las tortugas Chelidae (Pleurodira) y Testudinidae (Cryptodira) del Cenozoico argentino*. Tesis Doctoral, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, pp. 270.
- de la Fuente, M. 1992. Las tortugas Chelidae del Terciario superior y Cuaternario del territorio argentino. *Ameghiniana* 29: 211-229.
- Deynat, P. P. and Brito, P. M. 1994. Révision des tubercules cutanés de raies (Chondrichthyes, Batoidea) du bassin du Paraná, Tertiaire d'Amerique du Sud. *Annales de Paleontologie* 80: 237-251.
- Doering, A. 1882. *Informe oficial de la Comisión científica agregada al Estado Mayor General de la expedición al Río Negro*. Geología. Buenos Aires: 299-530.
- Domning, D.P. 1982. Evolution of manatees: a speculative history. *Journal of Paleontology* 56: 599-619.
- Donadío, O. 1983. Estado sistemático de los representantes fósiles del género *Tupinambis* en Argentina (Squamata, Sauria, Teiidae) y consideraciones paleambientales. *Boletín de la Asociación Herpetológica Argentina* 1: 8.
- Edmund, G. A. 1987. Evolution of the genus *Holmesina* (Pampatheriidae, Mammalia) in Florida, with remarks on the taxonomy and distribution. *Texas Memorial Museum, Pearce-Sellards Series* 45: 1-20.
- Emmons, L.H.. 1999. A New Genus and Species of Abrocomid Rodent from Perú (Rodentia: Abrocomidae). *American Museum Novitates* 3279: 1-14.
- Flynn, J.J. and Swisher, C.C. 1995. Cenozoic South American Land-mammal ages: correlation to global geochronologies. In: W.A. Berggren, D.V. Kent, and J. Handerbol (eds.). *Geochronology, Time scales, and Correlation: Framework for a Historical Geology*. SEPM Special Publication, 54: 317-333.
- Frailey, C.D. 1986. Late Miocene and Holocene mammals, exclusive of the Notoungulata, of the rio Acre region, western Amazonia. *Contribution on Science* 374: 1-46.
- Frenquelli, J. 1920a. Contribución al conocimiento de la geología de Entre Ríos. *Boletín de la Academia Nacional de Ciencias de Córdoba (Republica Argentina)* 24: 55-256.

- Frenguelli, J. 1920b. Apuntes sobre mamíferos fósiles entrerrianos. *Boletín de la Academia Nacional de Ciencias de Córdoba (República Argentina)* 24: 27-54.
- Frenguelli, G. 1922. Sulla contemporanea presenza del *Carcharodon megalodon* Ag. e del *Carcharodon rondeleti* M. et H. nel Paranense superiore de Entre Ríos (Rep. Argentina). *Bolletino de la Società Geologica Italiana* 41: 50-52.
- Frenguelli, G. 1930. Las guayquerías de San Carlos en la provincia de Mendoza. *Publicaciones del Departamento de Extensión Universitaria*, Universidad Nacional del Litoral 9: 1-54.
- Galván, A.F. and Ruiz Huidobro, O.J. 1965. Geología del valle de Santa María. Estratigrafía de las formaciones mesozoico-terciarias. *Actas de las Segundas Jornadas Geológicas Argentinas* Córdoba 3: 217-230.
- Gasparini, Z. 1996. Biogeographic Evolution of the South American Crocodylians. In: G. Arratia (ed.). *Contributions of Southern South America to Vertebrate Paleontology. München Geowissenschaftliche Abhandlungen* 30, pp. 159-184.
- Gasparini, Z., de la Fuente, M. and Donadio, O. 1986. Los reptiles cenozoicos de la Argentina: Implicancias paleoambientales y evolución biogeográfica. *Actas IV Congreso Argentino de Paleontología y Bioestratigrafía*, Mendoza 2: 119-130.
- Gayet, M. 1991. "Holostean" and Teleostean fishes of Bolivia. In: R. Suárez-Soruco (ed.). *Fósiles y facies de Bolivia. I. Vertebrados*. Revista Técnica de Yacimientos Petrolíferos Fiscales Bolivianos 12, pp. 453-494.
- Goin, F.J. 1991. Los Didelphoidea (Mammalia, Marsupialia) del Cenozoico tardío de la Región Pampeana. Unpublished thesis, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, 327 p.
- Goin, F.J. 1995. Los Marsupiales. En: M. T. Alberdi, G. Leone and E. P. Tonni (eds.). *Evolución biológica y climática de la Región Pampeana durante los últimos cinco millones de años. Un ensayo de correlación con el Mediterráneo Occidental*. Monografías, Museo Nacional de Ciencias Naturales y Consejo Superior de Investigaciones Científicas, Madrid, pp. 163-179.
- Goin, F.J. 1997a. New clues for the understanding of Neogene marsupial radiations in South America. In: R. F. Kay, R. H. Madden, R. L. Cifelli and J. Flynn (eds.). *Vertebrate Paleontology in the Neotropics. The Miocene Fauna of La Venta*, Colombia. Smithsonian Institution Press, Washington D. C., pp. 185-204
- Goin, F.J. 1997b. Sobre la edad y afinidades de *Zygolestes paranensis* Ameghino, 1898 (Marsupialia: Didelphidae: Marmosinae). *Neotrópica* 43: 15-19.
- Goin, F.J., Montalvo, C.I. and Visconti, G. (Submitted). Los Marsupiales (Mammalia) del Mioceno Superior de la Formación Cerro Azul (Provincia de La Pampa, Argentina). *Revista Española de Geología*, Madrid.
- Goin, F.J. and Pascual, R. 1987. News on the biology and taxonomy of the marsupials Thylascomilidae (Late Tertiary of Argentina). *Anales de la Academia Nacional de Ciencias Exactas, Físicas y Naturales* 39: 219-246.
- Gradstein, D. and Ogg, C. 1996. A geologic time scale. *Episodes* 19 (chart enclosed to this number).
- Iriondo, M.H. 1980. El cuaternario de Entre Ríos. Revista de la . *Asociación de Ciencias Naturales del Litoral* 11: 125-141.
- Iriondo, M.H. and E.D. Rodríguez. 1973. Algunas características sedimentológicas de la Formación Ituzaingó entre La Paz y Pueblo Brugo (Entre Ríos). *Actas del V Congreso Geológico Argentino* 1: 317-331.
- Kay, R.F., Madden, R.H., Vucetich, M.G., Carlini, A.A., Mazzoni, M.M., Re, G.H., Heizler, M. and Sandeman, H. 1999. Revised Age of the Casamayoran South American Land Mammal "Age". Climatic and Biotic Implications. *Proceedings of the National Academy of Sciences* 96: 13235-13240.
- Kraglievich, L. 1922. *Amphiocnus paranense* n. gen., n.sp. Un probable precursor del *Megalocnus* de la isla de Cuba en la formación Entrerriana. *Physis* 6: 73-87.
- Kraglievich, L. 1923. Descripción de dos cráneos y otros restos del género *Pliomorpus* Amegh. procedentes de la formación entrerriana de las Barrancas del Río Paraná. *Anales del Museo Nacional de Historia Natural de Buenos Aires* 33: 1-56.
- Kraglievich, L. 1931. Cuatro notas paleontológicas sobre *Octomyiodon averus* Amegh., *Argyrolagus palmeri* Amegh., *Tetrastylus montanus* Amegh. y *Muñizia paranensis* n. gen., n. sp. *Physis* 10: 242-266.
- Kraglievich, L. 1932. Diagnosis de nuevos géneros y especies de roedores cávidos y eumegámidos fósiles de Argentina. *Anales de la Sociedad Científica Argentina* 114: 155-181 and 211-237.
- Kraglievich, L. 1934. *La antigüedad pliocena de las faunas de Monte Hermoso y Chapadmalal, deducidas de su comparación con las que le precedieron y sucedieron*, pp. 1-136. Imprenta «El Siglo Ilustrado», San José 938, Montevideo.
- Kraglievich, J.L. 1952. El perfil geológico de Chapadmalal y Miramar, Provincia de Buenos Aires. *Revista Museo Municipal Ciencias Naturales y Tradicional Mar del Plata* 1:8-37.
- Langston, W. 1965. Fossil crocodylians from Colombia and the Cenozoic history of the Crocodylids in South America. *University of California Publications in Geological Sciences* 52: 1-157.
- Langston, W. and Gasparini, Z. 1997. Crocodylians, *Gryposuchus*, and the South American Gavials. In: R. Kay, R. Madden, R. Cifelli and J. Flynn (eds.), *Vertebrate Paleontology in the Neotropics. The Miocene Fauna of La Venta*. Smithsonian Institution Press 8: 113-154.
- Lapparent de Broin, Bocquentin, F.J., and Negri, F. 1993. Gigantic turtles (Pleurodira: Pelomedusidae) from the Late Miocene-Early Pliocene of south western Amazon. *Bulletin Institut française d'Études Andines* 22: 657-670.
- Larrazet, A. 1886. Des pièces de la peau de quelques Sélaciens fossiles. *Bulletin de la Société Géologique de France* 14:259-261.
- López, R. 1964. Problemas de la distribución geográfica de los peces marinos suramericanos. *Boletín del Instituto de Biología Marina* 7: 57-63.
- Lundberg, J. 1997. Fishes of the La Venta fauna: additional taxa, biotic and paleoenvironmental implications. In: R. F. Kay, R. H. Madden, R. L. Cifelli, and J. J. Flynn (eds.), *Vertebrate paleontology in the Neotropics: The Miocene fauna of La Venta*,

- Colombia. Smithsonian Institution Press, Washington, D.C, pp. 67-91.
- Lundberg, J., Machado-Allison, A. and Kay, R. 1986. Miocene characid fishes from Colombia: evolutionary stasis and extirpation. *Science* 234: 208-209.
- Lundberg, J.G., Linares, O.J., Antonio, M.E. and Nass, P. 1988. *Phractocephalus hemiliopterus* (Pimelodidae, Siluriformes) from the upper Miocene Urumaco Formation, Venezuela: a further case of evolutionary stasis and local extinction among South American fishes. *Journal of Vertebrate Paleontology* 8: 131-138.
- Marshall, L.G. 1979. Review of the Prothylacyninae, an extinct Sub family of South American "Dog-like" Marsupials. *Fieldiana Geology, New Series* 3: 1-50.
- Marshall, L.G. 1981. Review of the Hathyacyninae, an extinct subfamily of South American "Dog-like" Marsupials. *Fieldiana Geology, New Series* 7: 1-120.
- Marshall, L.G. 1985. Geochronology and Land-mammal biochronology of the transamerican faunal interchange. In: *The Great American Interchange*. D. Stehli and S.D. Webb. Plenum Press, p. 49-78.
- Marshall, L.G. 1987. Systematics of Itaboraian (Middle Paleocene) age "opossum-like" marsupials from the limestone quarry at São José de Itaboraí, Brasil. In: M. Archer (ed.), *Possums and Opossums, Studies in Evolution*. Surrey Beatty & Sons and The Royal Zool. Soc. New South Wales, Sydney, pp 91-160.
- Marshall, L.G. and Patterson, B. 1981. Geology and geochronology of the mammal-bearing Tertiary of the valle de Santa María and río Corral Quemado, Catamarca province, Argentina. *Fieldiana Geology* 9: 1-80.
- Marshall, L., Hoffstetter, R. and Pascual, R. 1983. Mammals and stratigraphy: geochronology of the continental mammal-bearing Tertiary of South America. *Palaovertebrata Mémoire Extraordinaire* 1-93.
- Marshall, L.G., Case, J.A. and Woodburne, M.O. 1990. Phylogenetic relationships of the families of Marsupials. In: H. H. Genoways (ed.), *Current Mammalogy*, Vol. 2. Plenum Press, New York, pp 433-505.
- Marshall, L.G., Sempere, T. and Gayet, M. 1993. The Petaca (Late Oligocene-Middle Miocene) and Yecua (Late Miocene) formations of the Subandean Chaco basin, Bolivia, and their tectonic significance. *Documents du Laboratoire de Géologie*, Lyon 125: 291-301.
- Marshall, L., Berta, A., Hoffstetter, R., Pascual, R., Reig, O., Bombin, M. and Mones, A. 1984. Mammals and stratigraphy: geochronology of the continental mammal-bearing Quaternary of South America. *Palaovertebrata Mémoire Extraordinaire* 1-76.
- Martin, L. 1983. The origin and early radiation of birds. In: A. H. Brush and G. A. Clark (eds.), *Perspectives in Ornithology, Essays presented for the Centennial of the American Ornithologists' Union*, Cambridge University Press, New York, pp. 291-338.
- Medem, F. 1983. *Los Crocodylia de Sur America. Volumen II*. Bogotá, Universidad Nacional de Colombia, pp.1-270.
- Moussy, M. de. 1857. Cuadro general de la Ciudad del Paraná. *Nacional Argentino*. 161, 162, 163, 164, Paraná.
- Moussy, M. de. 1860. *Description Physique, Géographique et Statistique de la Confédération Argentine*. I-III + Atlas.
- Mones, A. 1986. Palaovertebrata Sudamericana. Catálogo sistemático de América del Sur. Part 1: Lista preliminar y Bibliografía. *Courier Forschungsinstitut Senckenberg* 82: 1-50.
- Muizon, Ch. de and M. Bond. Les Phocidae (Mammalia) miocènes de la Formation Paraná (Entre Ríos, Argentine). *Bulletin du Muséum National d'Histoire Naturelle, Paris* 4: 165-207.
- Noriega, J.I. 1994. *Las Aves del "Mesopotamiense" de la provincia de Entre Ríos, Argentina*. Tesis Doctoral Nro. 611, Facultad de Ciencias Naturales, Universidad Nacional de La Plata, Argentina, 162 p.
- Noriega, J.I. 1995. The avifauna from the "Mesopotamian" (Ituzaingó Formation: Upper Miocene) of Entre Ríos Province, Argentina. *Courier Forschungsinstitut Senckenberg* 181: 141-148.
- Noriega, J.I. 1998. Aspectos paleozoogeográficos del registro de los Passeriformes (Aves) del Plioceno y Pleistoceno en la provincia de Buenos Aires. *V Jornadas Geológicas y Geofísicas Bonaerenses*, Mar del Plata 1: 65-71.
- Noriega, J.I. 2000. Nuevos restos de Phororhacidae (Aves: Gruiformes) del "Mesopotamiense" (Fm. Ituzaingó; Mioceno tardío) en la provincia de Entre Ríos, Argentina. *XVI Jornadas Argentinas de Paleontología de Vertebrados, San Luis*.
- Noriega, J.I. (In press). Body mass estimation and locomotion of the Miocene peleciform bird *Macranhinga*. *Acta Paleontologica Polonica*.
- Noriega, J.I. and Alvarenga, H. 2000. Phylogeny of the Tertiary giant darters (Pelecaniformes: Anhingidae) from South America. *5th International Meeting of the Society of Avian Paleontology and Evolution, Beijing*.
- North American Commission on Stratigraphic Nomenclature. 1983. North American Stratigraphic Code. *Bulletin American Association of Petroleum Geologists* 67: 841-875.
- Orbigny, A. d'. 1842. *Voyage dans l'Amérique méridionale*. P. Bertrand, Paris, 188 p.
- Pascual, R. and Odreman Rivas, O. 1971. Evolución de las comunidades de los vertebrados del Terciario argentino. Los aspectos paleozoogeográficos y paleoclimáticos relacionados. *Ameghiniana* 8: 372-412.
- Pascual, R. and Bondesio, P. 1972. Un roedor Cardiatheriinae (Hydrochoeridae) de la Edad Huayqueriense (Mioceno tardío) de La Pampa. Sumario de los ambientes terrestres en la Argentina durante el Mioceno. *Ameghiniana* 19: 19-35.
- Pascual, R., Vucetich, M.G. and Bond, M. 1985. Main pathways of mammalian diversification in South America. In: D. Stehli and S. D. Webb (eds.), *The Great American Biotic Interchange*. New York and London, Plenum Press, pp. 219-247.
- Pascual, R., Vucetich, M.G. and Scillato-Yané, G.J. 1990. Extinct and recent South American and Caribbean Megalonychidae

- edentates and Hystricognathi rodents: outstanding examples of isolation. International Symposium Biogeographical aspects of insularity, *Atti dei Convegni Lincei*, Roma 85: 627-640.
- Pascual, R., Ortega Hinojosa, E., Gondar, D. and Tonni E.P. 1965. Las edades del Cenozoico mamífero de la Argentina, con especial atención a aquellas del Territorio Bonaerense. *Anales de la Comisión de Investigación Científica* 6: 165-193.
- Perea, D. and Ubilla, M. 1989. Selacifauuna del Mioceno superior del Uruguay. *Boletín Sociedad Zoológica del Uruguay* 5: 11-12.
- Perea, D. and Ubilla, M. 1990. Los selacios (Chondrichthyes) de la Fm. Camacho (Mioceno sup., Uruguay). *Revista Sociedad Uruguaya de Geología* 4: 5-13.
- Perea, D.; Ubilla, M. and Martínez, S. 1985. Nuevos aportes a la fauna, geocronología y paleoambientes del Neoterciario del sur del Uruguay. *Boletín de la Sociedad Zoológica del Uruguay* 3: 42-54.
- Perea, D., Ubilla, M. and Piñeiro, G. 1996. First Fossil Record (Late Miocene) of *Phrynosops* (*P. geoffroanus* Complex: Chelidae) from Uruguay: Biostratigraphical and Paleoenvironmental Context. *Copeia* 2: 445-451.
- Pilleri, G. and Gühr, M. 1979. Skull, sonar field and swimming behaviour of *Ischyrorhynchus vanbenedeni* (Ameghino, 1891) and taxonomical position of the genera *Ischyrorhynchus*, *Anisodelphis* and *Pontoplanodes* (Cetacea). *Investigations on Cetacea* 10: 17-70.
- Priem, F. 1911. Poissons fossiles de la République Argentine. *Bulletin de la Société Belge de Géologie, Paléontologie et Hydrologie* 329-340.
- Rancy, A., Boquentin Villanueva, J., Pereira de Souza Filho, J., Santos, J.C.R. and Negri, F.R. 1989. Lista preliminar da fauna do Neógeno da região oriental do estado do Acre, Brasil (Material depositado em Rio Branco). VII Jornadas Argentinas de Paleontología de Vertebrados, Buenos Aires. *Ameghiniana* 26: 249.
- Rassmusen, D. and Kay, R. 1992. A Miocene aninga from Colombia, and comments of the zoogeographic relationships of South America Tertiary's avifauna. In: K. E Campbell (ed.). *Papers in Avian Paleontology honoring Pierce Brodkorb*. Natural History Museum of Los Angeles County, Science Series 36: 225- 230.
- Reig, O.A., 1957. Sobre la posición sistemática de "*Zygolestes paranensis*" Amegh. y de "*Zygolestes entrerrianus*" Amegh., con una reconsideración de la edad y correlación del "Mesopotamiense". *Holmbergia* 5: 209-226.
- Reig, O.A., 1958. Notas para una actualización del conocimiento de la fauna de la Formación Chapadmalal. II. Amphibia, Reptilia, Aves, Mammalia (Marsupialia: Didelphidae, Borhyaenidae). *Acta Geológica Lilloana* 2: 255-283.
- Reig, O. 1981. Teoría del origen y desarrollo de la fauna de mamíferos de América del Sur. *Monographie Naturae* 1: 1-1182.
- Reig, O.A., Kirsch, J.A.W. and Marshall, L.G. 1985. New conclusions on the relationships of the opossum-like marsupials, with an annotated classification of the Didelphimorphia. *Ameghiniana* 21: 335-343.
- Reig, O.A., Kirsch, J.A.W. and Marshall, L.G. 1987. Systematic relationships of the living and Neocenozoic American "Opossum-like" marsupials, with comments on the classifications of this and of the Cretaceous and Paleogene New World and European Metatherians. In: M. Archer (ed.). *Possums and Opossums, Studies in Evolution*. Surrey Beatty & Sons and The Royal Zoological Society of New South Wales, Sydney, pp. 1-89
- Reinhart, R.H. 1976. Fossil sirenians and demostylids from Florida and elsewhere. *Bulletin of the Florida State Museum, Biological Sciences* 20: 187-300.
- Rhodin, A.G. and Mittermeier, R.A. 1983. Description of *Phrynosops williamsi*, a new species of chelid turtle of the South American *P. geoffroanus* complex. In: A. G. Rhodin and K. Miyata (eds.), *Advances in herpetology and evolutionary biology*. Museum of Comparative Zoology, Harvard, pp. 58-73.
- Ridgway, S.H. and Harrison R. (eds). 1989. *Handbook of Marine Mammals: River Dolphins and the Larger Toothed Whales*. Academic Press, New York, 343 p.
- Riggs, E.S. and Patterson, B. 1939. Stratigraphy of Late-Miocene and Pliocene deposits of the Province of Catamarca (Argentina) with notes on the faunae. *Physis* 14: 143-162.
- Ringuélet, R.A. 1961. Rasgos fundamentales de la zoogeografía de la Argentina. *Physis* 22: 151-170.
- Ringuélet, R. 1975. Zoogeografía y ecología de los peces de aguas continentales de la Argentina y consideraciones sobre las áreas ictiológicas de América del Sur. *Ecotur* 2: 1-122.
- Ringuélet, R., Arámburu, R. and Alonso, A. 1967. *Los peces argentinos de agua dulce*. Comisión de Investigaciones Científicas de la Provincia de Buenos Aires, La Plata, 602 p.
- Río, C.J. del 1988. Bioestratigrafía y cronoestratigrafía de la Formación Puerto Madryn (Mioceno medio) Provincia del Chubut, Argentina. *Anales de la Academia Nacional de Ciencias Exactas, Físicas y Naturales* 40: 231-254.
- Río, C.J. del 1991. Revisión sistemática de los bivalvos de la Formación Paraná (Mioceno medio), provincia de Entre Ríos, Argentina. *Monografías de la Academia Nacional de Ciencias Exactas, Físicas y Naturales* 7: 11-90.
- Rusconi, C. 1933. Observaciones críticas sobre reptiles terciarios de Paraná (Alligatoridae). *Revista de la Universidad Nacional de Córdoba* 20: 57-106.
- Rusconi, C. 1934. Posición estratigráfica y relaciones zoológicas de los crocodylianos argentinos extinguidos. *Boletín Paleontológico* 4: 7-12.
- Rusconi, C. 1935. Observaciones sobre los gaviales fósiles argentinos. *Anales de la Sociedad Científica Argentina* 119: 203-214.
- Rusconi, C. 1948. El Puelchense de Buenos Aires y su fauna (Plioceno medio). (Primera parte). *Publicaciones del Instituto de Fisiografía y Geología, Universidad Nacional del Litoral* 33: 1-99.
- Rusconi, C. 1949. El Puelchense de Buenos Aires y su fauna (Plioceno medio). (Segunda parte). *Publicaciones del Instituto de Fisiografía y Geología, Universidad Nacional del Litoral* 36: 100-242.

- Sala, J.M. and Auge, M.P. 1970. Algunas consideraciones geohidrológicas del Noreste de la provincia de Buenos Aires. *Actas IV Jornadas Geológicas Argentinas* 2: 321-336.
- Santa Cruz, J.N. 1972. Estudio sedimentológico de la Formación Puelche en la Provincia de Buenos Aires. *Revista de la Asociación Geológica Argentina* 27: 5-62.
- Scartascini, G. 1954. El límite Plio-Pleistoceno de la Provincia de Entre Ríos. *Revista del Museo Municipal de Ciencias Naturales de Mar del Plata* 1: 55-77.
- Scillato-Yané, G.J. 1975a. Presencia de *Macroephractus retusus* (Xenarthra, Dasypodidae) en el Plioceno del Área Mesopotamia (Argentina). Su importancia bioestratigráfica y paleobiogeográfica. *Ameghiniana* 12: 322-328.
- Scillato-Yané, G.J. 1975b. Nuevo género de Dasypodidae (Edentata, Xenarthra) del Plioceno de Catamarca (Argentina). Algunas consideraciones filogenéticas y zoogeográficas sobre los Euphractini. *Actas del I Congreso Argentino de Paleontología y Bioestratigrafía*, Tucumán 2: 449-461, 1974.
- Scillato-Yané, G.J. 1977a. Octomylodontinae: una nueva Subfamilia de Mylodontidae (Edentata, Tardigrada). Descripción del cráneo y mandíbula de *Octomyodon robertoscagliai* n. sp. procedente de la Formación Arroyo Chasicó (Edad Chasicuense, Plioceno temprano) del sur de la Provincia de Buenos Aires (Argentina). Algunas consideraciones filogenéticas y sistemáticas sobre los Mylodontoidea. *Publicaciones del Museo Municipal de Ciencias Naturales. "L. Scaglia"* 2: 123-140.
- Scillato Yané, G.J. 1977b. Nuevo Megalonychidae (Edentata, Tardigrada) de Edad Chasicuense (Plioceno Temprano) del sur de la provincia de Buenos Aires (Argentina), su importancia filogenética, bioestratigráfica y paleobiogeográfica. *Revista de la Asociación de Ciencias Naturales del Litoral* 8: 45-54.
- Scillato-Yané, G.J. 1980. Nuevo Megalonychidae (Edentata, Tardigrada) del "Mesopotamiense" (Mioceno Tardío-Plioceno) de la provincia de Entre Ríos. *Ameghiniana* 17: 193-199.
- Scillato-Yané, G.J. 1981. Nuevo Mylodontinae (Edentata, Tardigrada) del "Mesopotamiense" (Mioceno Tardío-Plioceno) de la provincia de Entre Ríos. *Ameghiniana* 18: 29-34.
- Scillato-Yané, G.J. 1982. *Los Dasypodidae (Mammalia, Edentata) del Plioceno y Pleistoceno de Argentina*. Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, Tesis Doctoral: 1-159.
- Scillato-Yané, G.J. 1986. Los Xenarthra fósiles de Argentina (Mammalia, Edentata). *Actas II Congreso Argentino de Paleontología y Bioestratigrafía y I Congreso Latinoamericano de Paleontología*, Mendoza 2: 151-155.
- Simpson, G.G. 1940. Review of the mammal-bearing Tertiary of South America. *Proceedings of the American Philosophical Society* 83: 649-710.
- Soria, M.F. 1981. Los Litopterna del Colhuehuapense (Oligoceno tardío) de la Argentina. *Revista del Museo Argentino de Ciencias Naturales «Bernardino Rivadavia» e Instituto Nacional de Investigación de las Ciencias Naturales, Paleontología* 3: 1-54.
- Springer, S. and Bullis, H. R. 1960. A new species of sawshark, *Pristiophorus schroederi*, from the Bahamas. *Bulletin of the Marine Science of the Gulf and Caribbean* 10: 241-254.
- Tambussi, C.P. and Noriega, J.I. 1996. Summary of the avian fossil record from the southern South America. In: G. Arratia (ed.). *Contributions of Southern South America to Vertebrate Paleontology*. Münchner Geowissenschaftliche Abhandlungen 30, pp. 245-264.
- Tonni, E.P., Scillato-Yané, G.J., Cione, A.L. and Carlini, A. 1998. Bioestratigrafía del Mioceno continental en el curso inferior del arroyo Chasicó, provincia de Buenos Aires. In: *Resúmenes del VII Congreso Argentino de Paleontología y Bioestratigrafía*, Bahía Blanca, p. 135.
- Uliana, M.A. and Biddle, K.T. 1988. Mesozoic-Cenozoic paleogeographic and geodynamic evolution of southern South America. *Revista Brasileira de Geociências* 18: 172-190.
- Verzi, D.H., Vucetich, M.G. and Montalvo, C.I. 1995. Un nuevo Eumysopinae (Rodentia, Echimyidae) del Miceno tardío de la Provincia de la Pampa y consideraciones sobre la historia de la subfamilia. *Ameghiniana* 3: 191-195.
- Vizcaíno S.F. and Scillato-Yané G.J. 1995. An Eocene tardigrade (Mammalia, Xenarthra) from Seymour Island, West Antarctica. *Antarctic Science* 7: 407-408.
- Vucetich, M.G. and Verzi, D.H. 1995. Los Roedores Caviomorpha. In: M. T. Alberdi, G. Leone, and E. P. Tonni (eds.). Evolución biológica y climática de la Región Pampeana durante los últimos cinco millones de años. *Monografías del Museo de Ciencias Naturales de Madrid* 12: 211-225.
- Vucetich, M.G. and Verzi, D.H. 1996. Un nuevo Eumysopinae (Rodentia, Echimyidae) de la "Formación" Irene (Chapadmalalense inferior?) y la diversidad de la Subfamilia. *Actas IV Jornadas de Geología y Geofísica Bonaerenses* Tomo 1: 159-165.
- Vucetich, M.G., Verzi, D.H. and Tonni, E.P. 1997. Paleoclimatic implications of the presence of *Clyomys* (Rodentia, Caviomorpha) in the Upper Pliocene of Buenos Aires Province. *Palaogeography, Palaoclimatology, Palaecology* 128: 207-214.
- Wieland, G. 1923. A new Paraná pleurodiran. *American Journal of Science* 5: 1-14.
- Windhausen, A. 1931. *Geología Argentina*. Segunda Parte: Geología Histórica y Regional del territorio Argentino, Jacobo Peuser Lda. Editores, Buenos Aires, pp. 1-645.
- Wood, R.C. 1976. Two new species of *Chelus* from the Late Tertiary of northern South America. *Breviora* 435: 1-26.
- Wood, R.C. 1997. Turtles. In: R. Kay, R. Madden, R. Cifelli and J. Flynn (eds.). *Vertebrate Paleontology in the Neotropics: The*

- Miocene Fauna of La Venta*. Smithsonian Institution Press, pp. 155-170.
- Wood, R.C. and Díaz de Gamero, M.L. 1971. *Podocnemis venezuelensis*, a new fossil pelomedusid (Testudines, Pleurodira) from the Pliocene of Venezuela and a review of the history of *Podocnemis* in South America. *Breviora* 376: 1-23.
- Woodward, A.S. 1900. On some fish-remains from the Parana Formation, Argentine Republic. *Annals and Magazine of Natural History*, London 6: 1-7.

Appendix

Vertebrates not included in the tables
Marine taxa

Chondrichthyans

Order Squaliformes
Family Squalidae
Squalus sp.
Order Lamniformes
Family Odontaspidiidae
Carcharias acutissima
Family Lamnidae
Isurus hastalis
Family "Otodontidae"
Carcharocles megalodon,
Order Carchariniformes
Family Carcharhinidae
Carcharhinus spp.
Galeocerdo aduncus
Family Hemigaleidae
Hemipristis serra,
Order Heterodontiformes
Family Heterodontidae
Heterodontus sp.
Order Squatiniformes
Family Squatinidae
Squatina sp.
Order Myliobatiformes
Family Dasyatidae
Family Myliobatidae

«Osteichthyans»

Order Perciformes
Family Sciaenidae
Sciaenidae indet.

Cetaceans

Odontoceti
Family Pontoporiidae
Pontistes rectifrons Burmeister, 1891
Family Physeteridae
Physeteridae indet.

Mysticeti
Family Balaenopteridae
Balaenopteridae indet.
Family Balaenidae
Balaenidae indet.

Sirenians

Familia Dugongidae

?*Dioplotherium* sp.

Aquatic carnivores

Family Phocidae

Properiptychus argentinus (Ameghino, 1893)

Continental taxa

CHONDRICHTHYANS

Order Myliobatiformes

Family Potamotrygonidae

«Osteichthyans»

Order Siluriformes

Family Pimelodidae

Pimelodidae indet.

Family Loricariidae

Loricariidae indet.

Family Callichthyidae

Callichthyidae indet.

Family Ariidae

Ariidae indet.

Family Auchenipteridae

Auchenipteridae indet.

Order Characiformes

Family Characidae

Colossoma macropomus

Family Cynodontidae

Cynodontidae indet.

«Reptilians»

Family Chelidae

Phrynos sp.

Family Testudinidae

Testudinine indet.

Family Alligatoridae

Caiman jacare (Daudin), 1802

Caiman latirostris (Daudin), 1802

Caiman australis (Burmeister), 1885

Family Nettosuchidae

Mourasuchus nativus (Gasparini), 1985

Family Gavialidae

Gryposuchus neogaeus (Rusconi), 1935

Family Teiidae

Tupinambis cf. *T. meriana* (Dumeril & Bibron) 1839

Birds

Order Pelecaniformes

Family Anhingidae

Macranhinga paranensis Noriega 1992

Family Anhingidae

Indeterminate genus and species

Order Charadriiformes

Family Palaelodidae

Megapalaelodus sp.
 Family Phoenicopteridae
 Phoenicopteridae indet.
 Order Anseriformes
 Family Anatidae
 Subfamily Dendrocheninae
 Dendrocheninae indet.
 Order Ciconiiformes
 Family Ciconiidae
 Tribu Mycteriini
 Mycteriini indet.
 Order Rheiformes
 Family Rheidae
 Rheidae indet.
 Order Gruiformes
 Family Rallidae
 Rallidae indet.
 Family Phororhacidae
 Subfamily Phororhacinae
Onactornis? pozzi Kraglievich 1931
 Subfamily Palaeociconiinae
Andalgalornis steulleti Kraglievich 1931
Andalgalornis deautieri Kraglievich 1931

Marsupials

Sparassodonta
 Family Hathliacynidae
Notictis ortizi Ameghino 1889
 Family Borhyaenidae
 Subfamily Prothylacyninae
Stylocynus paranensis Mercerat 1917
 Family Thylacosmilidae
Achlysictis lelongi Ameghino 1891
 Didelphimorphia
 Family Didelphidae
 Subfamily Marmosinae
Zygolestes paranensis Ameghino 1898

Cetaceans

Odontoceti
 Family Iniidae
Ischyrorhynchus vanbenedeni Ameghino, 1891
Saurocetes argentinus Burmeister, 1871
Saurocetes gigas (Cozzuol, 1989)
 Family Pontoporiidae
 Pontoporiinae? indet.

Sirenians

Family Trichechidae
Ribodon limbatus Ameghino, 1883

Terrestrial carnivores

Family Procyonidae

Cyonasua argentina

Recibido: 18 de agosto de 2000

Aceptado: 15 de noviembre de 2000