




Asociación
Parasitológica
Argentina

Volumen 13. Nro. 2

Órgano oficial de difusión científica de la Asociación Parasitológica Argentina

(Rev. Arg. Parasitol.)



ISSN: 2313-9862

Revista Argentina de Parasitología

Eyes wide open: First record of eyeworms of genus *Oxyspirura* (Nematoda-Thelaziidae) in Argentina

Con los ojos bien abiertos: Primer registro de gusanos oculares del género *Oxyspirura* (Nematoda-Thelaziidae) en Argentina

Drago Fabiana B.^{1,2*}, Amigo Martina^{1,3}, Podesta Taiel^{1,3}, Draghi Regina¹, Núñez Verónica^{1,2}

ABSTRACT: The eyeworm, *Oxyspirura octopapillata* (Nematoda-Thelaziidae) is reported for the first time in Argentina parasitizing the eyes of the little nightjar, *Setopagis parvula* (Aves-Caprimulgidae). The host was collected in Formosa Province, Argentina, and nine specimens belonging to this nematode species were found. Its morphological characteristics and morphometric data are described, which include a divided buccal capsule, 17 male caudal papillae and a spicular ratio of 1: 2.3-2.9. These features allow us to differentiate these specimens from the other Neotropical species of the genus *Oxyspirura*. This is the first parasitological study of *S. parvula* in Argentina.

Keywords: Nematodes, *Oxyspirura octopapillata*, *Setopagis parvula*

RESUMEN: El gusano ocular, *Oxyspirura octopapillata* (Nematoda-Thelaziidae) es citado por primera vez en Argentina parasitando los ojos del atajacaminos chico, *Setopagis parvula* (Aves-Caprimulgidae). El hospedador fue colectado en la provincia de Formosa, Argentina y se encontraron nueve especímenes pertenecientes a esta especie de nematode. Se describen sus características morfológicas y sus datos morfométricos, que incluyen una cápsula bucal dividida, 17 papilas caudales en el macho y una relación espicular de 1: 2.3-2.9. Estas características permiten diferenciar a los ejemplares del presente estudio de las otras especies Neotropicales del género *Oxyspirura*. Este es el primer estudio parasitológico de *S. parvula* in Argentina.

Palabras clave: Nematodes, *Oxyspirura octopapillata*, *Setopagis parvula*

The little nightjar, *Setopagis parvula* (Gould) (Caprimulgiformes: Caprimulgidae) is an exclusively Neotropical bird. Its helminth fauna is scarcely known, to date it was only reported as host of three helminth species: one nematode *Oxyspirura petrowi* Skrjabin, 1929 (Thelaziidae) from Brazil, and two cestodes *Metadilepis spasskiorum* Georgiev and Vaucher, 2003 and *Mariauxilepis paraguayensis* Georgiev and Vaucher, 2003 (Metadilepididae) from Paraguay (Vicente *et al.*, 1995; Georgiev and Vaucher, 2003). The goal of this work was to increase the knowledge of the parasitic helminths of *S. parvula*.

One specimen of *S. parvula* was hunted with a shotgun in September 2012 at La Marcela farm, Pirané, Formosa Province, Argentina (26° 17' 35" S; 59° 06' 38" W). The authorization was provided by Ministerio de la Producción y Ambiente of Formosa Province (N° 003516). The bird was kept frozen at

-20°C, then thawed, dissected, and examined under a stereoscopic microscope in the laboratory. The parasitological examination revealed the presence of nine nematodes belonging to the genus *Oxyspirura* Drasche, 1897, found on the eyes surface, under the nictitating membranes. Helminths were removed and preserved in 70% ethanol. For examination, specimens were cleared by immersion in Amman's lactophenol. Measurements are given in millimeters (mm) unless otherwise stated, expressed as a range. The photographs were taken with a camera phone (Xiaomi MiA3). Both nematodes and the host were deposited in the Helminthological and Ornithological Collections of the Museo de La Plata, La Plata, Argentina (MLP- He 8132 and MLP-O-P 14812, respectively). Identification of nematodes was carried out following the keys proposed by Ybarra (1948), Chabaud (1975), and Oliveira-Rodrigues (1978).

¹División Zoología Invertebrados, Museo de La Plata, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata (UNLP). Paseo del Bosque S/N°, 1900 La Plata, Buenos Aires, Argentina.

²Comisión de Investigaciones Científicas de la provincia de Buenos Aires (CIC).

³Facultad de Ciencias Naturales y Museo, UNLP.

Nematoda Rudolphi, 1808

Family Thelaziidae Skrjabin, 1915

Oxyspirura octopapillata Caballero, 1942 (Tables 1-2, Figs. 1-2)

Description based on four males and five females. Nematodes with approximately the same width along the body, slightly attenuated at anterior end and pointed at posterior end. Males slightly shorter than females, with a spiral pointed tail. Males and females exhibited cervical alae extending approximately to the level of the nerve ring. Mouth terminal. Buccal capsule divided into an anterior part wider (0.028-0.052) than the posterior one (0.014-0.023). Esophagus slightly divided into anterior muscular and posterior glandular portions. Muscular portion 0.11-0.23 long in males and 0.2-0.31 in females. Glandular portion 0.86-0.92 long in males and 0.89-1.18 in females. Excretory pore not observed.

Males with two unequal and dissimilar spicules, both transversely striated, being the right spicule shortest, straight and robust, and the left spicule as the longest, curved and thin. Average spicular length ratio of 1: 2.6. Gubernaculum weakly sclerotized observed in only one male. Caudal papillae arranged in nine precloacal symmetrical pairs, and eight postcloacal pairs, with the first three pairs slightly asymmetrical and the remaining five symmetrical.

In the Neotropical realm, 19 species of *Oxyspirura* are known parasitizing avian hosts (Jairapuri and Siddiqi, 1967; Baruš, 1968; Guerrero, 1969; Oliveira-Rodrigues, 1978; Vicente *et al.*, 1995). Among them, eight species have a buccal capsule divided into an anterior part wider than the posterior one, like the specimens here studied: *O. cephaloptera* (Molin, 1860) Stossich, 1897 reported parasitizing Coraciiformes

and Passeriformes from Brazil (Oliveira-Rodrigues, 1978), *O. chauvancyi* Díaz Ungría, 1963 described parasitizing Passeriformes from French Guiana (Díaz Ungría, 1963), *O. diazungriai* Guerrero, 1969 and *O. guriensis* Guerrero, 1969 in Galliformes and Coraciiformes, respectively from Venezuela (Guerrero, 1969), the cosmopolitan species *O. mansoni* (Cobbold, 1879) Ransom, 1904 reported in numerous wild and domestic birds (Oliveira-Rodrigues, 1978), *O. navali* Caballero, 1936 described parasitizing Accipitriformes from Mexico (Caballero, 1936), *O. octopapillata* Caballero, 1942 reported parasitizing Accipitriformes, Falconiformes and Caprimulgiformes from Brazil, Mexico and Cuba (Caballero, 1942; Ybarra, 1948; Baruš, 1968; Vicente *et al.*, 1995), and *O. tanasijtchuki* (Skrjabin, 1916) Oliveira-Rodrigues, 1964 reported in Passeriformes from Paraguay (Cram, 1927).

The specimens here studied can be differentiated from seven of the eight species previously mentioned in the Neotropical realm by the spicular ratio and number of caudal papillae in males. *Oxyspirura cephaloptera* possesses a spicular ratio of 1: 5, seven precloacal pairs and six postcloacal pairs of caudal papillae (Oliveira-Rodrigues, 1978). *Oxyspirura chauvancyi* has a spicular ratio of 1: 13, three precloacal pairs, one adcloacal pair, and two postcloacal pairs of caudal papillae (Díaz-Ungría, 1963). *Oxyspirura diazungriai* possesses a spicular ratio of 1: 14.5, three precloacal pairs, one adcloacal pair, and two postcloacal pairs of caudal papillae (Guerrero, 1969). *Oxyspirura guriensis* has a spicular ratio of 1: 15.5, three precloacal pairs, one adcloacal pair, and two postcloacal pairs of caudal papillae (Guerrero, 1969). *Oxyspirura navali* possesses a spicular ratio of 1: 1.85, two precloacal pairs, and six postcloacal pairs of caudal papillae (Caballero, 1936).




Figure 1. Female of *Oxyspirura octopapillata*. a) Dorsal view of anterior end showing buccal capsule (bc), nervous ring (nr) and cervical alae (ca). b) Lateral view of anterior end showing muscular esophagus (me) and glandular esophagus (ge). c) Lateral view of posterior end showing vulva (v) and anus (a). Scale bars 100 μ .




Figure 2. Posterior end of *Oxyspirura octopapillata* male. a) Right spicule and precloacal papillae. The arrows indicate the precloacal papillae. b) Left spicule. Scale bars 50 μ .

Oxyspirura tanasijtchuki has a spicular ratio similar to specimens here studied (1: 2.4), but possesses five precloacal papillae (2 symmetrical pairs and one fifth isolated papilla) and seven postcloacal caudal papillae (1 symmetrical pair and 5 asymmetrical papillae) (Cram, 1927). *Oxyspirura mansonii* possesses a spicular ratio of 1: 15, four precloacal pairs, one adcloacal pair, and four postcloacal pairs of caudal papillae (Guerrero, 1969, Oliveira-Rodrigues, 1978).

The specimens here studied have a similar spicular ratio and number of caudal papillae that the eighth species, *O. octopapillata*. Also, the specimens recovered from *S. parvula* possess morphological and morphometric characters similar to those described

by Caballero (1942), Ybarra (1948), and Baruš (1969) for this species (Tables 1-2).

Eyeworms belonging to the genus *Oxyspirura* are heteroxenous nematodes found in a wide variety of wild and domestic birds. These nematodes are usually found on the eye surface, under the nictitating membrane, as well as in the lacrimal ducts and other eye glands. The adults of *Oxyspirura* spp. deposit the eggs, which together with lacrimal secretions follow the tear ducts to the mouth where they are swallowed and eliminated through the feces. The eggs are ingested by cockroaches, crickets or grasshoppers that act as intermediate hosts (Anderson, 2000; Kalyanasundaram et al., 2019).

Table 1. Comparative morphometric data for females of *Oxyspirura octopapillata* in different hosts and localities. Measurements in mm, unless otherwise stated.

Source	Present study	Caballero (1942)	Ybarra (1948)	Baruš (1969)
Country	Argentina	Mexico	Mexico	Cuba
Hosts	<i>Setopagis parvula</i> (Gould) (Caprimulgidae)	<i>Caracara plancus cheriway</i> (Jacquin) (Falconidae)*	<i>Nyctidromus albicollis</i> (Gmelin) (Caprimulgidae)	<i>Caracara plancus audubonii</i> (Cassin) (Falconidae)**
Body length	15.1-18	18.9	17.5-18	14.77
Body width	0.400-0.484	0.520	0.380	0.480
Buccal capsule depth	0.040-0.053	0.041	0.045	0.039
Buccal capsule width	0.038-0.042	0.037	0.038	0.035
Nerve ring (d _{fae})	0.22-0.37	0.266	0.220	0.269
Esophagus total length	1.2-1.375	1.36	1.4	1.35
Muscular esophagus width	0.048-0.067	0.049	0.046	0.058
Glandular esophagus width	0.100-0.154	0.072	0.165	0.160
Vulva (d _{fpe})	0.532-0.992	1.140	0.980	0.949
Anus (d _{fpe})	0.107-0.410	0.440	0.460	0.409
Eggs (μ)	40-58 x 20-33	53-57 x 29-33	49-53 x 30-32	40-46 x 25-27

*cited as *Polyborus cheriway*, **cited as *Polyborus cheriway audubonii*. Abbreviations: d_{fae}- distance from anterior end; d_{fpe}- distance from posterior end.

Table 2. Comparative morphometric data for males of *Oxyspirura octopapillata* in different hosts and localities. Measurements in mm.

Source	Present study	Caballero (1942)	Ybarra (1948)
Country	Argentina	Mexico	Mexico
Hosts	<i>Setopagis parvula</i> (Gould) (Caprimulgidae)	<i>Caracara plancus cheriway</i> (Jacquin) (Falconidae)*	<i>Nyctidromus albicollis</i> (Gmelin) (Caprimulgidae)
Body length	11.7-13.6	13.54	15.8-16.5
Body width	0.330-0.420	0.400	0.300-0.340
Buccal capsule depth	0.036-0.048	0.037**	0.045
Buccal capsule width	0.028-0.052	0.029	0.035
Nerve ring (d _{fae})	0.100-0.257	0.217	0.250-0.260
Esophagus total length	0.970-1.15	1.266	1.35
Muscular esophagus width	0.043-0.055	0.053	0.057
Glandular esophagus width	0.106-0.120	0.164	0.180-0.190
Left spicule length	0.310-0.490	0.462	0.418
Right spicule length	0.110-0.190	0.195	0.205
Spicular length ratio	1: 2.30-1: 2.90	1: 2.75	1: 2.04
Caudal papillae	9 pairs precloacal 8 pairs postcloacal	9 pairs precloacal 8 pairs postcloacal	9 pairs precloacal 1 pair adcloacal 7 pairs postcloacal

*cited as *Polyborus cheriway*. **erroneously reported as 0.37 mm. Abbreviations: d_{fae}- distance from anterior end.

Eyeworms, such as *O. petrowi*, negatively impact ocular tissues by causing inflammation of the lacrimal ducts, keratitis and lesions on the Harderian glands. In time, the adenitis would likely result in gland atrophy and fibrosis. These conditions cause a deficiency in tear production called keratoconjunctivitis sicca, which is commonly known as “dry eye”. This condition could cause corneal lacerations, reduced vision and decreased survival (Dunham *et al.*, 2016).

This is the first report of the genus *Oxyspirura* in Argentina, and the first parasitological study of *S. parvula* in Argentina.

The authors express their gratitude to Francisco and Carlos Montoya during their stay in Formosa Province. We also thank Agustín Abba and Luis Pagano for their assistance in collecting the host.

This work was supported by Universidad Nacional de La Plata (11/N979).

LITERATURE CITED

Anderson, R. C. (2000). Nematode Parasites of Vertebrates: Their Development and Transmission. 2nd Edition. Wallingford, Oxford, United Kingdom: CAB International.

Baruš, V. (1968). Parasitic nematodes of birds of the family Icteridae (Passeriformes) in Cuba. Folia Parasitologica, 15, 131-146.

Baruš, V. (1969). Nematodes Parasitic in birds of Cuba. Acta societatis zoologicae Bohemoslovaca, 33, 193-210.

Caballero y C., E. (1936). Contribución al conocimiento de los nematodos de las aves de México III. Anales del Instituto de Biología de la Universidad Nacional Autónoma de México, 7, 469-475.

Caballero y C. E. (1942). Nematodos de las aves de México IX. Descripción de una nueva especie del género *Oxyspirura* y consideraciones acerca de las especies mexicanas ya conocidas. Anales del Instituto de Biología, Serie Zoológica, 13, 527-532.

Cram, E. B. (1927). Bird parasites of the nematode suborders Strongylata, Ascaridata and Spirurata. Bulletin United States National Museum, 140, 1-465.

Chabaud A. G. (1975). Keys to genera of the order Spirurida. Part. I. Camallanoidea, Dracunculoidea, Gnathostomatoidea, Physalopteroidea, Rictularioidea and Thelazioidea. In: R. C. Anderson, A. G. Chabaud, and S. Willmott (Eds.). CIH Keys to the Nematode Parasites of Vertebrates (334-360). Farnham Royal: Commonwealth Agricultural Bureaux.

Díaz-Ungría, C. (1963). Nematodes parásitos colectados por la misión Chauvancy en Guyana Francesa. Bulletin du Muséum National d’Histoire Naturelle, 2° Série, 4, 441-453.

Dunham, N. R., Reed, S., Rollins, D. and Kendall, R. J. (2016). *Oxyspirura petrowi* infection leads to pathological consequences in Northern bobwhite (*Colinus virginianus*). International Journal for

- Parasitology: Parasites and Wildlife, 5, 273-276.
- Georgiev, B. B. and Vaucher, C. (2003). Revision of the Metadilepididae (Cestoda: Cyclophyllidea) from Caprimulgiformes (Aves). *Revue Suisse de Zoologie* 110 (3): 491-532.
- Guerrero, R. (1969). Contribución al conocimiento del género *Oxyspirura* Drasche in Stossich, 1897, con descripción de dos especies nuevas (Nematoda, Spirurida, Thelaziidae). *Memoria de la Sociedad de Ciencias Naturales La Salle*, 29 (82), 84-101.
- Jairapuri, D. S. and Siddiqi, A. H. (1967). A Review of the Genus *Oxyspirura* Drasche in Stossich, 1879 (Nematoda: Thelaziidae) with descriptions of fourteen new species. *Journal of Helminthology*, 4, 337-363.
- Kalyanasundaram, A., Brym, M. Z., Blanchard, K. R., Henry, C., Skinner, K., Henry, B. J., Herzog, J., Hay, A. and Kendall, R. J. (2019). Life-cycle of *Oxyspirura petrowi* (Spirurida: Thelaziidae), an eyeworm of the northern bobwhite quail (*Colinus virginianus*). *Parasites & Vectors*, 12, 555.
- Oliveira-Rodrigues, H. (1978). Estudo das espécies da subfamilia Oxyspirurinae Yamaguti, 1961 referidas para o Brasil (Nematoda, Spiruroidea) (Tese de Mestrado). UFRJ.
- Vicente, J. J., Oliveira Rodrigues, H., Corrêa Gomes, D. and Magalhães Pinto, R. (1995). Nematóides do Brasil. Parte IV: Nematóides de Aves. *Revista Brasileira de Zoologia* 12 (Supl. 1), 1-273.
- Ybarra, G. A. (1948). Estudio monográfico de nematodos parásitos de las aves de México (Tesis). Universidad Nacional Autónoma de México.

Recibido: 29 de Agosto de 2024

Aceptado: 30 de septiembre de 2024
