

Toppan Best-set Premedia Limited	
Journal Code: JZO	Proofreader: Mony
Article No: JZO886	Delivery date: 13 December 2011
Page Extent: 12	

1 **Body mass estimation in Early Miocene native South American ungulates: a predictive equation based on 3D landmarks**

4 **G. H. Cassini^{1,2}, S. F. Vizcaíno³ & M. S. Bargo³**

1 División Mastozoología, Museo Argentino de Ciencias Naturales 'Bernardino Rivadavia', CONICET, Ciudad Autónoma de Buenos Aires, Argentina
2 Departamento de Ciencias Básicas, Universidad Nacional de Luján, Luján, Buenos Aires, Argentina
3 División Paleontología Vertebrados, Museo de La Plata, La Plata, Argentina

Keywords

Notoungulata; Litopterna; Astrapotheria; Patagonia; Miocene; Santacrucian; body mass; centroid size.

Correspondence

Guillermo H. Cassini, División Mastozoología, Museo Argentino de Ciencias Naturales 'Bernardino Rivadavia', Consejo Nacional de Investigaciones Científicas y Técnicas, Av. Angel Gallardo 470, Ciudad Autónoma de Buenos Aires, C1405DJR, Argentina. Tel: +54 011 4982 0306, ext. 208
Email: gcassini@macn.gov.ar

3 Editor: ••••

Received 17 August 2011; revised 14 November 2011; accepted 24 November 2011

doi:10.1111/j.1469-7998.2011.00886.x

Introduction

5 South American native ungulates include several extinct taxa that evolved within the context of South American geographic isolation during most of the Cenozoic (Bond, 1999). The ungulates of the Santacrucian Age (Early Miocene), belonging to the orders Notoungulata, Litopterna and Astrapotheria, are particularly interesting for paleobiological studies due to their diversity, richness and quality of specimen preservation (Cassini *et al.*, in press). Their biological diversity includes types that resemble equids, rhinos, camels and caviomorph rodents in both shape and size (see Cassini & Vizcaíno, in press), as well as a giant form with no analogue in the current faunas.

6 A recent study of Santacrucian ungulates from south-east Patagonia (Cassini *et al.*, in press) provides an extensive description of their diversity in a bounded geographic and stratigraphic context. Notoungulates are represented by four genera of Typotheria: *Interatherium* and *Protypotherium*

Abstract

South American native ungulates include extinct taxa that evolved within the geographical context given by the isolation of South America during most of the Cenozoic. The ungulates (orders Notoungulata, Litopterna and Astrapotheria) of the Santa Cruz Formation (late Early Miocene) are particularly interesting for paleobiological studies due to their diversity, richness and quality of preservation of the specimens. The body mass estimation of extinct species is one of the basic biological attributes for paleobiological reconstructions. The most common way to estimate body mass from fossils is using linear regression. Here, we used geometric morphometric techniques in order to estimate their body mass. We used regressions based on centroid size of 3D craniomandibular landmark configurations, including extant ungulates (their size and ecological relatives). Cases were weighted to maximize the taxonomic evenness. A broad body size range was recorded. The highest predictive power is obtained with those functions derived from the highest taxonomic and ecological diversity. The highest taxonomic richness corresponds to masses below 100 kg. Among Notoungulata, typotheres (*Hegetotheriidae* + *Interatheriidae*) vary from 1 to less than 10 kg, while the smaller toxodontid reached 100 kg and the larger 500 kg. Litoptern proterotheriid body masses vary from 10 to 50 kg, and macraucheniids surpass 100 kg. The astrapotheres (*Astrapotheria*) reached (or even surpassed) 1000 kg, being the only megamammal in the Santacrucian ungulate assemblage.

(*Interatheriidae*), and *Hegetotherium* and *Pachyrukhos* (*Hegetotheriidae*). Notoungulata also comprise three genera of Toxodontia: *Nesodon* and *Adinotherium* (*Toxodontidae*), and *Homalodotherium* (*Homalodotheriidae*). The litopterns include representatives of the families Proterotheriidae (*Anisolophus*; *Diadiaphorus*; *Tetramerorhinus*; *Thoatherium*) and Macraucheniidae (*Theosodon*). Astrapotheres are represented by a single genus (*Astrapotherium*).

Toxodontids are medium- to large-sized notoungulates often compared with rhinoceroses both in appearance and ecological role. They have a long and high skull, high crowned cheek teeth, with lateral incisors developed as tusks, and a robust skeleton. Typotheres are small- to medium-sized mammals, mostly described as rodent-like in overall form, although *Hegetotheriidae* more closely resemble hares (Reguero, Dozo & Cerdeño, 2007) and *Interatheriidae* resemble a composite of a small mustelid's body with a hyrax's head (Croft & Anderson, 2008; Cassini *et al.*, in press). Typotheres have wide and low skulls, ever growing high crowned cheek

teeth, and gracile postcranial skeletons. Among Litopterna, macraucheniiids include large animals with some convergences with camelids (long neck and a canal for the vertebral artery in cervicals, which passes through the neural arch; Scott, 1910). Proterotheriids are medium-sized animals, mostly described as similar to small primitive holarctic horses due to their body size and overall cranial shape, but the presence of reduced digits and mesaxonic limbs makes convergent with modern equids. Their tooth crown morphology, on the other hand, resembles that of artiodactyls such as deer (Bond *et al.*, 2001). Astrapotheres are large to very large mammals (including strict megamammals, i.e. 1000 kg or more *sensu*; Owen-Smith, 1988), described as morphologically intermediate between a tapir and an elephant, with large canine tusks and a putative proboscis (Scott, 1928; Riggs, 1935).

Size is one of the most variable features in the diversity of Santacrucian ungulates. Usually measured through body mass (bm), size is one of the most important aspects of the biology because it is highly correlated with various aspects of life such as diet, locomotion, energy, ecology, morphology and physiology (see Brown & West, 2000 and references therein).

Since the initial formalization of the allometric relationship by Kleiber (1932), various subsequent studies have applied the concept of allometry in physiological to ecological contexts (Calder, 1996). These studies demonstrated that many fundamental characteristics of biological organisms scale with corporal size following a power law formalized mathematically as $Y = aX^b$ (Savage *et al.*, 2004).

Body mass constitutes one of the first parameters to evaluate in the fossil taxa for the purpose of predicting biological attributes that are not preserved, and to propose paleoecological hypotheses (Andrews, Lord & Evans, 1979; Reed, 1998; Vizcaíno *et al.*, 2006). Although some attempts have been made to estimate the bm of the Santacrucian ungulates, the results are either restricted to few taxa or based only on a small set of measures. For example, Villafañe (2005) reported bm's of proterotheriids, and Scarano, Carlini & Illius (2011) for the species of *Protypotherium*. Croft (2000) and Vizcaíno *et al.* (2010) used a more inclusive sample of taxa; their estimations were based on the first lower molar dimensions and the mean of a set of skull measurements, respectively.

Predicting bm of extinct species using allometric functions is the most common approach (e.g. Fariña, Vizcaíno & Bargo, 1998; Millien & Bovy, 2010). Some authors postulated that some postcranial elements (e.g. humerus and femur) are more accurate estimators of bm because they provide body support (Damuth & MacFadden, 1990), especially for species lacking modern representatives or falling outside the size ranges of extant species. Thus, postcrania have been extensively used as bm proxies (e.g. Alexander *et al.*, 1979; Scott, 1990; Farlow *et al.*, 2005). However, in the mammalian fossil record, cranial remains are more frequently recovered and much easier to assign taxonomically and therefore are frequently used as bm estimators as well (e.g. Gingerich, Smith & Rossenberg, 1982; Myers, 2001; Mendoza, Janis & Palmqvist, 2006).

In geometric morphometric analyses, shape and size are decomposed and centroid size (*cs*) is used as a proxy for size

(Goodall, 1991; Dryden & Mardia, 1998). In the absence of allometry, *cs* is not correlated with shape (Bookstein, 1986; Kendall, 1986). Hood (2000) evaluated the potential use of *cs* in the study of size sexual dimorphism in the muskrat *Ondatra zibethicus* (Rodentia: Muridae, Arvicolinae) as well as eight bat species (Chiroptera, Pteropodidae). He concluded that *cs* is a geometric measure of size that follows the same mathematical behaviour as bm, indicating that *cs* is an excellent variable indicative of bm. Meloro & O'Higgins (2011) showed that mandibular *cs* is highly correlated with mean body weight in Carnivora. Ercoli & Prevosti (2011) obtained significant log-linear regressions between bm and *cs* for appendicular skeleton of extant carnivorous marsupials and placental mammals, and used *cs* to estimate the bm of species of Santacrucian Sparassodonta metatherians.

In this contribution, we attempt to obtain equations to predict bm from *cs* of three-dimensional (3D) landmark configurations of cranium and mandible using a broad taxonomical and ecological diversity of extant ungulates.

Materials and methods

Data

Fifty-nine specimens of ungulates with little to no apparent deformation from the Patagonian Santa Cruz Formation (late Early Miocene) were analyzed: 19 toxodontids and 24 typotheres (Notoungulata), 11 proterotheriids and 1 macraucheniid (Litopterna), and 3 astrapotheres (Astrapotheria) (see Appendix SI – online supporting information). 7

The reference sample of extant ungulates consisted of 505 individuals belonging to 155 species (134 Artiodactyla, 16 Perissodactyla and 5 Hyracoidea; Appendix SII – online supporting information). All extant families and subfamilies were represented (Table 1). Only adult specimens with third molar erupted and in occlusion were included, and captive animals were avoided as much as possible.

Landmark data

3D landmark coordinates were acquired with a Microscribe G2L digitizer; they are defined in Tables 2 and 3 and are shown in Fig. 1a–c. Both sides and midline landmarks were included. These comprise 36 cranial (see Milne & O'Higgins, 2002) and 14 mandibular landmarks including type I (or anatomic) and II (or mathematic). 8

Regression functions

All statistics, mathematical operations, regression functions and prediction of mass were carried out in R.

Museum specimens rarely include body weight; therefore, analyses were performed using the mean bm values reported in the literature (see Appendix SII). The reference sample includes species with sexual dimorphism; when information on male and female bm data was available, they were differentiated. We also obtained equations using the mean bm by

Table 1 Number of species/genera sampled in relation to the total number of species/genera in each family/subfamily

Taxa	Species sampled	Total of species	% species	Genera sampled	Total of genera	% genera
Perissodactyla						
Equidae	8	8	100%	1	1	100%
Tapiridae	4	4	100%	1	1	100%
Rhinocerotidae	4	5	80%	3	4	75%
Artiodactyla						
Suidae	9	19	47%	5	5	100%
Tayassuidae	3	3	100%	3	3	100%
Hippopotamidae	2	2	100%	2	2	100%
Camelidae	6	6	100%	3	3	100%
Ruminantia						
Tragulidae	3	8	38%	3	3	100%
Giraffidae	2	2	100%	2	2	100%
Antilocapridae	1	1	100%	1	1	100%
Moschidae	2	7	29%	1	1	100%
Cervidae						
Capreolinae	13	22	59%	8	9	89%
Hydropotinae	0	1	0%	0	1	0%
Muntiacini	3	12	25%	2	2	100%
Cervini	10	16	63%	6	7	86%
Bovidae						
Bovinae	10	24	42%	6	9	67%
Reduncinae	8	9	89%	3	3	100%
Antilopinae	15	38	39%	10	15	67%
Caprinae	25	35	71%	11	12	92%
Aepycerotinae	1	1	100%	1	1	100%
Cephalophinae	9	18	50%	3	3	100%
Hippotraginae	6	8	75%	3	3	100%
Alcelaphinae	6	10	60%	4	4	100%
Hyracoidea						
Procaviidae	5	7	71%	3	3	100%
Total	155	266	58%	85	98	87%

species. Centroid size (cs) of landmark configuration (M) is a size function defined as the square root of the sum of squared distances from each landmark to the centroid of the configuration and is independent of position or orientation of the configuration (Claude, 2008). The cs of each specimen was calculated using this R command: $cs \leftarrow \sqrt{\text{sum}(\text{apply}(M, 2, \text{var})) * (\text{dim}(M)[1] - 1)}$ (modified from Claude, 2008).

Body mass (bm) grows at cube rates, while cs does so linearly, so the two variables were log(base 10) transformed, converting the relationship of these variables from a power formula ($bm = a cs^b$) to a straight line: $\log(bm) = \log(a) + b \log(cs)$. Variable normality was analyzed with a Shapiro–Wilk test and normal quantile–quantile plot on log-transformed variables (Shapiro & Wilk, 1965). We assessed the extent and significance of the association between cs and bm via Spearman rank non-parametric correlations. We used the original non-transformed data because it tests how monotonically they are related, even if their relationship is not linear (Zar, 1999).

The regression functions were fitted to the data by means of ordinary least squares (OLS) models on log10-transformed variables. The OLS regression model is recommended if the main aim of our regression analysis is prediction (Warton *et al.*,

2006). Additionally, quantile regression analysis was performed. It was developed by Koenker & Bassett (1978) as a new class of robust alternatives to the OLS regression. This method is suitable for estimating functional relations between variables for all portions of a probability distribution (Cade & Noon, 2003). Quantile regression estimates can be used to construct prediction and tolerance intervals without assuming any parametric error distribution. Also, the estimates retain their statistical properties under any linear or non-linear monotonic transformation (e.g. logarithmic) as a consequence of this ordering property (Koenker & Machado, 1999). This allows the back transformation of the bm estimates to the original scale without any loss of information (Cade & Noon, 2003).

Some groups are widely represented in the database, while others (e.g. Procaviidae) are under-represented. The number of taxonomic groups is limited by the available sample (see Table 1), but their taxonomic evenness can be maximized (for an extensive discussion, see Mendoza *et al.*, 2006). In order to avoid biases, the taxa were weighted so each taxon contributes equally in the bm function; that is, in a family with one genus with four species and another with only two, both genera contribute in the same way to the equation. For this purpose, a

Table 2 Cranial landmarks, names and definitions used in the present study (see also Milne & O'Higgins, 2002)

Number	Name	Definition of landmark
1 & 22	<i>Stephanion</i>	<i>Sutura coronalis</i> at the intersection with the <i>linea temporalis</i>
2 & 23		<i>Sutura frontolacrimalis</i> at the <i>margo orbitalis</i>
3 & 24	<i>Dacryon</i>	<i>Sutura zygomaticolacrimalis</i> at the <i>margo orbitalis</i>
4 & 25		Anterior point of the <i>processus zygomaticus</i> of the <i>os temporale</i>
6 & 27		Superior root of the <i>zygoma</i> (on the <i>temporal bone</i>)
7 & 28		<i>Sutura zygomaticotemporal</i> on the <i>margo ventrale</i> of the <i>arcus zygomaticus</i>
8 & 29		Anteriormost origin of <i>musculus masseter</i> /Lowest point of the 'masseteric process'
9 & 30		Distal <i>margo alveolare</i> of the last molar
10 & 31		Mesial <i>margo alveolare</i> of first functional premolar
11 & 32		<i>Foramen infraorbitale</i>
12 & 33		<i>Sutura maxilloinsciva</i> at the <i>margo intralveolaris</i>
13 & 34		Distal <i>margo alveolare</i> of the lateralmost incisor or horny pad scar
14 & 35		<i>Sutura nasoinsciva</i> at the margin of the nasal aperture
15 & 36		<i>os nasale, frontale</i> and <i>maxilla</i> junction
16	Lambda	<i>Sutura occipitoparietalis</i>
17	Bregma	<i>Sutura coronalis</i> in the midline
18	<i>Nasion</i>	<i>Sutura frontonasalis</i> on the sagittal plane
19	<i>Rhinion</i>	Rostral edge of the <i>os nasale</i> on the <i>sutura internasalis</i>
20	<i>Nasospinale</i>	<i>Sutura interincisiva</i> on the mid-sagittal plane of nasal aperture
21	<i>Prosthion</i>	<i>Sutura interincisiva</i> on the <i>margo alveolaris</i>

new weighing variable was obtained following two approaches: one developed by Mendoza *et al.* (2006) and the other by De Esteban-Trivigno, Mendoza & De Renzi (2008). This variable is implemented as a 'weight' argument of 'lm' R function.

When predicted *bm* values from a least squares regression are retransformed back to arithmetic units, log-adjustment could yield biased estimates (Smith, 1993). In order to correct this potential bias, the ratio estimator (RE) (Snowdon, 1991) was applied. This ratio consisted of the mean of the observed values divided by the mean of retransformed predicted values. The predicted masses for the South American ungulates were retransformed back and then multiplied by the corresponding RE.

Although the correlation coefficient is one of the commonest ways to evaluate regressions, it poorly indicates the independent variable predictive power because it is affected by the range of values of the dependent variable and, therefore, it could be high even with high residuals (Smith, 1984). Thus, function reliability was evaluated by directly using their

Table 3 Mandibular landmarks, names and definitions used in the present study

Number	Name	Definition of landmarks
1	<i>Gnathion</i>	Caudo-ventral margin of the mandibular symphysis on the midline
2	<i>Infradentale</i>	<i>Alveoli dentalis</i> of i1 in the midline
3		Distal margin of <i>alveoli dentalis</i> of the last lower incisor
4		Distal margin of <i>alveoli dentalis</i> of the lower canine
5		Mesial margin of the <i>alveoli dentale</i> of the first functional premolar
6		<i>Margo interalveolaris</i> between last premolar and first molar
7		Distal margin of the <i>alveoli dentale</i> of the last molar
8	<i>Coronion</i>	The most superior point on the <i>processus coronoideus</i>
9	Mandibular notch	The most inferior point on the <i>incisura mandibulae</i>
10	<i>Condylion medial</i>	Most medial margin of mandibular condyle
11	<i>Condylion lateral</i>	Most lateral margin of mandibular condyle
12		Most dorsal-caudal rugosity from <i>musculus masseter</i> insertion
13		Most anterior roughness from <i>musculus masseter</i> insertion
14		<i>Foramen mandibulae</i>

percent prediction error as $\%PE = 100 \times [\text{observed} - \text{predicted}] / \text{predicted}$ (Smith, 1984).

Results

Figure 2 shows the box and whisker plot diagram of both variables *bm* and *cs*. They are discriminated between families across the three orders of extant ungulates and subfamilies among Bovidae and Cervidae. The *cs* values in these taxonomic groups show a pattern similar to *bm* (Fig. 2). Both log-transformed variables show no significant deviations from normality. Spearman's rank non-parametric correlation shows that cranial and mandibular *cs* are highly correlated with mean *bm* in the extant ungulates (cranial: $n = 155$, $r_s = 0.968$, $P < 0.0001$; mandibular: $n = 147$, $r_s = 0.976$, $P < 0.0001$).

All regressions obtained were statistically significant (Table 4). The *bm* range for all regressions is from 1.46 kg (represented by the bush hyrax *Heterohyrax brucei*) to 1750 kg (represented by the javan rhino *Rhinoceros sondaicus*). All determination coefficients (R^2) are high (Table 4) and the mean prediction error (%PE) is low. The relationship between the two variables, both for cranium and mandible, was isometrical [i.e. slope values around three expected under geometric similarity between a variable that grows linearly (*cs*) and cube (*bm*); Peters, 1983]. Based on %PE, we chose the equations for cranium and mandible obtained using the Mendoza *et al.* (2006) weighted approach (i.e. Cr.4 and Md.4 equations).




Figure 1 Landmark configuration on *Adinotherium* cranium (a, b) and mandible (c). See Tables 2 and 3 for their definition. The landmarks are shown on the right side and midline.

Only for two cranial regressions outliers were found. The outlier species are listed in Table 5 and eliminated from dataset for the calculation of regressions.

The distribution of extant taxa around regression line is shown in Fig. 3. Those groups with proportionately large skulls (consequently high *cs*) relative to their *bm* lie below the regression line (among others, suids, equids, and some alcelaphine bovids). On the other hand, some taxa with proportionately short skulls (e.g. tapirs, rhinos and saiga) lie above the regression line. However, they are not outliers.

Figure 4 presents a summary of quantile regression results for Cr.4 regression. The horizontal axis represents the quantile

scale and the vertical indicates the estimates. The conditional median and mean fits for both estimates (intercept and slope) are very similar as both 90% confidence intervals overlap. Also, the quantile intercepts and slopes between 0.15 and 0.7 remain enclosed to OLS estimate confidence intervals. Note that one consequence of this is that the OLS fit provides a very good estimate of the conditional mean in the sample. However, departures of the OLS model are represented by the larger than 0.7 quantile.

Estimated *bm* of each specimen of Santacrucian ungulates are listed on Appendix SI. Additionally, the minimum and maximum *bm* based on quantile regression are shown.

Mean *bm* at species level are listed in Table 6, discriminated between cranial and mandibular equations. Also, mean max *bm* are shown. The values obtained for these two elements are quite similar, except for *Astrapotherium magnum* (higher *bm* from mandible) and *Nesodon imbricatus* (lower *bm* from mandible). Except for typotheres, mandible mean max *bm* are always greater than cranial ones. For less-abundant species, the value for a single specimen is presented.

Among the assemblage of Santacrucian ungulates, *Astrapotherium* reaches higher *bm* (c. 700 kg). Among Notoungulata, *Nesodon* is the largest within Toxodontia (c. 400 kg), while both *Adinotherium* species are below 100 kg. All Typotheria are smaller than 10 kg; among them, *Protypotherium australe* and *Hegetotherium mirabile* are the largest (4.6 and 3.6 kg, respectively), and *Interatherium* spp. and *Pachyrukhos moyani* are the smallest (just below of 2 kg). Among Litopterna, all the Proterotheriidae fall in the 10–60 kg range. Only the Macraucheniidae *Theosodon* reaches more than 110 kg, being the largest Santacrucian litoptern.

Discussion

All regression equations have similar parameters (Table 4), indicating minor differences between the weighted and unweighted methods. In agreement with Mendoza *et al.* (2006) and De Esteban-Trivigno *et al.* (2008), this suggests that high taxonomical diversity leads to good predictive power functions. The equations obtained here have very low %PE values (6%) when compared with simple regressions that use skull or dental measurements of different mammalian lineages (16–97 %PE extracted from the literature; De Esteban-Trivigno *et al.*, 2008: table 3). Our results match the smallest %PE of multiple regressions previously reported (4.5–7.5%; Mendoza *et al.*, 2006: table 2). Although working with fragmentary fossils using equations based on a single variable (e.g. Janis, 1990) is convenient for simplicity, with almost complete specimens, our model predictions are more accurate due to smaller prediction errors and absence of allometry.

The distribution of taxa around the regression line indicates that if the taxon whose *bm* we attempt to estimate has a large skull relative to the body, the *bm* obtained could be overestimated. On the contrary, if they have a typical skull size, the OLS regression provides a reliable estimate.

The fitted values of the superior quantile regression return the maximum mean *bm* for a given *cs*. There is a low probability that an ungulate reaches a *bm* higher than that esti-




Figure 2 Box plot diagram (median, first and third quartile, and range) of body mass (top) and centroid size (bottom) within families and subfamilies categories of extant mammal sample.

mated by superior quantile regression. Therefore, we conclude that the superior quantile regression establishes a ‘theoretical’ maximum *bm* for a given centroid size.

Body mass estimation of extinct species

Applying a regression equation based on the first molar length (Damuth, 1990), Croft (2000) obtained the first *bm* estimations for Santacrucian ungulates. Croft’s values for every taxon are lower than every mean value obtained here. On the contrary, *bm* estimations, which are based on a mean of several equations taken from Janis (1990), obtained by Vizcaino *et al.* (2010) are higher (Table 6). Differences between cranial and mandibular *bm* estimates of *Astrapotherium* would

reflect lower cranial *cs* due to retraction of rostral elements and a higher mandible *cs* due to a long diastema.

Most recent *bm* estimations for notoungulates (Reguero, Candela & Cassini, 2010; Elissamburu, 2011; Scarano *et al.*, 2011) provide *bm* estimates larger than those obtained here (Table 6), although our *bm* of *Pachyrukhos moyani* (1.56 kg) is very similar to the value of 1.77 kg obtained by Elissamburu (2011).

The *bm* values obtained by the superior quantile regression should be interpreted as the maximum mean *bm* reached by an ungulate of a given *cs*. For some taxa, they are very close to the values reported in the aforementioned works. Among proterotheriids, quantile max *bm* values for *Anisolophus* and *Diadiaphorus* were very similar to those of Vizcaino *et al.*

Table 4 Statistical summary of regressions obtained using the log₁₀(centroid size) as independent variable and log₁₀(body mass) in kg as dependent variable

Regressions	<i>n</i>	<i>R</i> ²	Intercept	Std. error	<i>t</i> _(<i>n</i>-2)	Slope	Std. error	<i>t</i> _(<i>n</i>-2)	RE	%PE
Sex mean										
Cr.1	238 (145 sp)	0.953	-6.506	0.120	-54.116	3.083	0.044	69.775	1.084	6.554
Md.1	230 (140 sp)	0.955	-5.665	0.109	-52.174	2.980	0.043	69.391	1.086	6.442
Species mean										
Cr.2	155	0.950	-6.539	0.155	-42.128	3.093	0.057	54.153	1.108	6.392
Md.2	147	0.956	-5.720	0.135	-42.449	3.000	0.053	56.129	1.093	6.287
A. Weighted species mean										
Cr.3	155	0.950	-6.786	0.160	-42.383	3.186	0.059	53.746	1.044	6.552
Md.3	147	0.953	-5.926	0.143	-41.525	3.085	0.057	54.134	1.033	6.748
B. Weighted species mean										
Cr.4	150	0.976	-6.701	0.113	-59.416	3.165	0.041	76.989	1.000	6.0996
Qsup			-6.926	0.178	-38.888	3.321	0.070	47.143		
Qinf			-5.902	0.385	-15.316	2.763	0.129	21.339		
Md.4	147	0.965	-5.921	0.124	-47.854	3.088	0.049	63.435	1.001	6.0604
Qsup			-6.316	0.136	-46.277	3.345	0.074	45.057		
Qinf			-5.477	0.201	-27.278	2.826	0.082	34.463		

A, weighted as in De Esteban-Trivigno *et al.* (2008); B, weighted as in Mendoza *et al.* (2006); *n*, number of observations (if not indicated correspond also to species number); Qsup, superior quantile regression (0.99); Qinf, inferior quantile regression (0.01); %PE, percent prediction error; RE, correction factor; *t*_(*n*-2), Student's *t*-test value for both parameters. All were significant at the 0.05 level (*P* < 0.00001).

Table 5 Excluded data from indicated regression equation

Regression	Species	<i>r</i> -Student	<i>P</i> -value	<i>P</i> -Bonf
Cr.1	<i>Phacochoerus africanus</i> (M)	3.802	0.00018	0.044
Cr.4	<i>Catagonus wagneri</i>	5.161	<0.0001	<0.001
Cr.4	<i>Antilocapra americana</i>	4.646	0.00001	0.001
Cr.4	<i>Tayassu pecari</i>	4.469	0.00002	0.002
Cr.4	<i>Pecari tajacu</i>	5.435	<0.0001	<0.001
Cr.4	<i>Phacochoerus africanus</i>	4.742	<0.0001	0.001

P-value, unadjusted; *P*-Bonf, Bonferroni adjusted *P*-value. Critical value of *r*-Student equals 2.

(2010), but far lower than those of Villafañe (2005). Similarly, among notoungulates, for the toxodonts, *Adinotherium* and *Nesodon*, cranial max *bm* were similar to those obtained by Vizcaino *et al.* (2010) and Elissamburu (2011). While in typotheres, with the exception of *Pachyrukhos moyani*, all max *bm* values were lower than previously reported.

In summary, the OLS *bm* estimates lie between those based on dental and craniomandibular measurements, suggesting that estimations based on *cs* functions could be interpreted as a 'consensus' between them. This is supported by the fact that craniomandibular measurement estimates are in the range of our 'theoretical' maximum *bm*'s obtained from quantile regression.

Paleoecology


Although the Santacrucian Age span for over several million years (~22–14 Ma, Blisniuk *et al.*, 2005), all the genera and all the size range considered in this study have been recorded in a single locality (Puesto La Costa; Tauber, 1997; Vizcaino *et al.*,

2010; Cassini *et al.*, in press), in fossiliferous levels that may represent just different facies of the same environment.

The coexistence of many herbivores in a living assemblage implies partitioning of vegetation as a food source (Hirst, 1975). Size is one attribute of the niche differentiation in herbivorous mammals (Jarman & Sinclair, 1979; Owen-Smith, 1988): particularly among African bovids, the small-sized forms are more selective than the larger ones (Jarman & Sinclair, 1979). Also, food quality is inversely proportional to the fiber content (Illius, 1997), and while larger ungulates fed mainly on grasses, smaller ones do it on dicots (Gordon & Illius, 1996).

The Santacrucian ungulates can be separated into three size groups of base-10 logarithmic scale ranges, each log unit corresponding to one order of magnitude. The small-sized ungulates (1–10 kg) are exclusively typotheres; the medium-sized ungulates (10–100 kg) are the litoptern proterotheriids and the toxodontids *Adinotherium* spp.; and the large-sized ungulates (100–1000 kg) are the macraucheniid *Theosodon garretorum*, the toxodontid *Nesodon imbricatus* and the astrapother *Astrapotherium magnum*.

According to the previous statements, we should expect typotheres to be highly selective dicot feeders, proterotheriids dicot selective feeders, *Adinotherium* spp. and macraucheniiids mixed feeder, and *Nesodon* and astrapotheres grass bulk feeders. This scheme fits only partially with recent ecomorphologic and biomechanic feeding characterizations (see Cassini *et al.*, 2011; Cassini & Vizcaino, in press). Typotheres and *Astrapotherium* depart from this pattern, the former being highly grazers and the latter a selective feeder. As described earlier, both have morphologies much different from extant ungulates toxodontids and litopterns. Typotheres, in particular, are usually described as rodent-like. Perhaps like rodents



1 **Figure 3** Dispersion diagram of cranial regression Cr.4 (a) and mandibular regression Md.4 (b). Solid line: ordinary least squares regression; dashed line:
2 superior quantile regression; broken line: inferior quantile regression; asterisks: outliers removed from equation. Taxa symbols: cross – Hyacoidea;
3 squares – Perissodactyla (black – Equidae; gray – Tapiridae; white – Rhinocerotidae); white triangle – Hippopotamidae; black triangle – Suiformes; black
4 inverted triangle – Tragulidae; white inverted triangle – Antilocapridae; black rhombus – Camelidae; circle – Bovidae; gray discs – Cervidae.




Figure 4 Ordinary least squares and quantile regression estimates for **Cr.4** model. Solid line: ordinary least squares (OLS) estimate; dotted lines: 90% confidence intervals of OLS estimates. Shaded gray area: 90% pointwise confidence band for the quantile regression estimates.

Table 6 Mean body mass (kg) of Santacrucian ungulates obtained by the regressions equations and those previously reported in the literature

Species	Cranium			Mandible			Croft (2000)	Other sources	Vizcaino et al. (2010)
	Mean \pm SD	Max	n	Media \pm SD	Max	n			
<i>Astrapotherium magnum</i>	704.08 \pm 260.18	1,241.98	2	656.13 \pm 213.75	1,416.22	3	503.8		1021.63
<i>Theosodon garrettorum</i>				113.09 \pm 0.00	210.14	1	46.04 ^a		
<i>Anisolophus australis</i>				13.13 \pm 0.00	20.40	1	8.84 ^b	62.63(a)	18.14
<i>Diadiaphorus majusculus</i>	47.49 \pm 1.19	73.23	4	56.08 \pm 1.54	98.31	2	26.48	190.11(a)	70.25
<i>Tetramerorhinus cingulatum</i>	27.08 \pm 0.59	40.61	2					82.76(a)	
<i>Tetramerorhinus mixtum</i>	19.33 \pm 0.00	28.52	1					65.58(a)	
<i>Thoatherium minusculum</i>	19.79 \pm 0.00	29.23	1	15.16 \pm 0.00	23.83	1	6.25	45.00(a)	21.0
<i>Hegetotherium mirabile</i>	4.67 \pm 0.70	6.42	3				2.190	9.69 ^b (b)	14.23
<i>Hegetotherium mirabile</i>	1.56 \pm 0.00	2.04	1					1.77 ^b (b)	
<i>Interatherium robustum</i>	1.81 \pm 0.43	2.37	5	1.62 \pm 0.36	2.12	6	0.399	3.33 ^b (b)	3.5
<i>Interatherium excavatum</i>	1.40 \pm 0.00	1.82	1						
<i>Interatherium extensum</i>	1.33 \pm 0.00	1.72	1						
<i>Protypotherium australe</i>	3.63 \pm 0.36	4.93	5	3.82 \pm 0.00	5.35	1	2.817	6.74 ^b (b)/ 5.86(c)	7.8
<i>Protypotherium attenuatum</i>	1.93 \pm 0.00	2.54	1	1.84 \pm 0.00	2.43	1	0.906	2.85(c)	4.4
<i>Protypotherium praerutilum</i>	2.38 \pm 0.10	3.16	3				0.976	4.57(c)	
<i>Adinotherium ovinum</i>	58.40 \pm 11.67	91.07	9				33.75	119.45 ^b (b)	121.0 ^b
<i>Adinotherium robustum</i>	82.06 \pm 6.46	130.00	2				24.32		
<i>Nesodon imbricatus</i>	320.25 \pm 38.34	542.59	3	429.67 \pm 88.13	893.72	7	293.66	587.9 ^b (b)	554.0 ^b

^aBody mass obtained for *Theosodon lydekkeri*.

^bWhen body mass was reported only for the genus.

Other sources: (a) Villafañe (2005); (b) Elissamburu (2011); (c) Scarano et al. (2011).

SD, standard deviation.

(Langer, 2002) and unlike ungulates (Pérez-Barbería & Gordon, 2001) in typotheres, fiber content was not correlated with bm. This issue remains as a working hypothesis for future works.

Acknowledgements

We thank the Dirección de Patrimonio Cultural (Santa Cruz, Argentina); Dr Reguero (MLP), Dr Kramarz (MACN), Dr

Flynn (AMNH) and Dr Joyce (YPM) for access to vertebrate paleontological collections; Dr Verzi (MLP), Dr Flores (MACN), Dr Simmons (AMNH) and Dr Zyskowski (YPM) for access to mammalogy collections; Dr Toledo for the illustrations; and Dr Haber and the three anonymous reviewers for their valuable suggestions that improved the paper. The study of the YPM collections was partially funded by the John H. Ostrom Research Fund to GHC. This is a contribution to the Projects PICT 0143, PIP 1054 and UNLP N647 to S. F. V.

References

- Alexander, R.M., Jayes, S.A., Maloiy, G.M.O. & Wathuta, M.E. (1979). Allometry of the limb bones of mammals from shrews (*Sorex*) to elephant (*Loxodonta*). *J. Zool. (Lond.)* **189**, 305–314.
- Andrews, P., Lord, J.M. & Evans, E.M.N. (1979). Patterns of ecological diversity in fossil and modern mammalian faunas. *Biol. J. Linn. Soc.* **11**, 177–205.
- Blisniuk, P.M., Stern, L.A., Chamberlain, C.P., Idleman, B. & Zeitler, K.P. (2005). Climatic and ecologic changes during Miocene surface uplift in the Southern Patagonian Andes. *Earth Planet. Sci. Lett.* **230**, 125–142.
- Bond, M. (1999). Quaternary native ungulates of Southern South America. A synthesis. In *Quaternary of South America and Antarctic Peninsula: 177–205*. Rabassa, J. & Salemme, M. (Eds). Ushuaia, Tierra de Fuego: Centro Austral de Investigaciones Científicas and Universidad Nacional de la Patagonia.
- Bond, M., Perea, D., Ubilla, M. & Tauber, A.A. (2001). *Neolicanophrium recens* Frenguelli, 1921, the only surviving Proterotheriidae (Litopterna, Mammalia) into the South American Pleistocene. *Palaeovertebrata* **30**, 37–50.
- Bookstein, F.L. (1986). Size and shape spaces for landmark data in two dimensions: comment. *Stat. Sci.* **1**, 181–222.
- Brown, J.H. & West, G.B. (2000). *Scaling in biology*. New York: Oxford University Press.
- Cade, B.S. & Noon, B.R. (2003). A gentle introduction to quantile regression for ecologists. *Front. Ecol. Environ.* **1**, 412–420.
- Calder, W.A. (1996). *Size, function and life history*. Cambridge: Harvard University Press.
- Cassini, G.H., Cerdeño, E., Villafañe, A.L. & Muñoz, N. (in press). Paleobiology of Santacrucian native ungulates (Meridiungulata: Astrapotheria, Litopterna and Notoungulata). In *Early Miocene paleobiology in Patagonia: high-latitude paleocommunities of the Santa Cruz Formation: ••••*. Vizcaíno, S.F., Kay, R.F. & Bargo, M.S. (Eds). Cambridge: Cambridge University Press.
- Cassini, G.H., Mendoza, M., Vizcaíno, S.F. & Bargo, M.S. (2011). Inferring habitat and feeding behaviour of early Miocene notoungulates from Patagonia. *Lethaia* **44**, 153–165.
- Cassini, G.H. & Vizcaíno, S.F. (in press). An approach to the biomechanics of the masticatory apparatus of early Miocene (Santacrucian Age) South American ungulates (Astrapotheria, Litopterna, and Notoungulata): moment arm estimation based on 3D landmarks. *J. Mammal. Evol.* ••, ••••.
- Claude, J. (2008). *Morphometrics with R*. New York: Springer.
- Croft, D.A. (2000). *Archaeohyracidae (Mammalia: Notoungulata) from the Tinguiririca Fauna, central Chile, and the evolution and paleoecology of South American mammalian herbivores*. PhD thesis, University of Chicago at Chicago.
- Croft, D.A. & Anderson, L.C. (2008). Locomotion in the extinct notoungulate *Protypotherium*. *Palaeontol. Electron.* **11**, 1–20.
- Damuth, J. (1990). Problems in estimating body masses of archaic ungulates using dental measurements. In *Body size in mammalian paleobiology: estimation and biological implications: 229–253*. Damuth, J. & MacFadden, B.J. (Eds). Cambridge: Cambridge University Press.
- Damuth, J. & MacFadden, B.J. (1990). Introduction: body size and its estimation. In *Body size in mammalian paleobiology: estimation and biological implications: 1–10*. Damuth, J. & MacFadden, B.J. (Eds). Cambridge: Cambridge University Press.
- De Esteban-Trivigno, S., Mendoza, M. & De Renzi, M. (2008). Body mass estimation in Xenarthra: a predictive equation suitable for all quadrupedal terrestrial placentals? *J. Morphol.* **269**, 1276–1293.
- Dryden, I.L. & Mardia, K.V. (1998). *Statistical shape analysis*. Chichester: John Wiley & Sons.
- Elissamburu, A. (2011). Estimación de la masa corporal en géneros del Orden Notoungulata. *Estud. Geol. (Madr)* ••, ••••. doi: 10.3989/egol.40336.133.
- Ercoli, M.D. & Prevosti, F.J. (2011). Estimación de masa de las especies de Sparassodontata (Mammalia, Metatheria) de la edad santacrucense (Mioceno temprano) a partir del tamaño del centroide de los elementos apendiculares: inferencias paleoecológicas. *Ameghiniana* **48**, 605–620.
- Fariña, R.A., Vizcaíno, S.F. & Bargo, M.S. (1998). Body mass estimations in Lujanian (Late Pleistocene-Early Holocene of South America) mammal megafauna. *Mastozoology. Neotrop.* **5**, 87–108.
- Farlow, J.O., Hurlburt, G.R., Else, R.M., Britton, A.R.C. & Langston, W. Jr (2005). Femoral dimensions and body size of *Alligator mississippiensis*: estimating the size of extinct mesoeucrocodylians. *J. Vertebr. Paleontol.* **25**, 354–369.
- Gingerich, P.D., Smith, B.H. & Rossenberg, K. (1982). Allometric scaling in the dentition of primates and prediction of body weight from tooth size in fossils. *Am. J. Phys. Anthropol.* **58**, 81–100.
- Goodall, C.R. (1991). Procrustes methods in the statistical analysis of shape. *J. R. Stat. Soc. Ser. B (Methodol.)* **53**, 285–339.
- Gordon, I.J. & Illius, A.W. (1996). The nutritional ecology of African Ruminants: a reinterpretation. *J. Anim. Ecol.* **65**, 18–28.
- Hirst, S.M. (1975). Ungulate habitat relationships in a South African woodland savanna ecosystem. *Wildl. Monogr.* **44**, 1–60.
- Hood, C.S. (2000). Geometric morphometric approaches to the study of sexual size dimorphism in mammals. *Hystrix* **11**, 77–90.
- Illius, A.W. (1997). Physiological adaptation in savanna ungulates. *Proc. Nutr. Soc.* **56**, 1041–1048.
- Janis, C.M. (1990). Correlation of cranial and dental variables with body size in ungulates and macropodoids. In *Body size*

- 1 *in mammalian paleobiology: estimation and biological impli-*
2 *cations: 255–300.* Damuth, J. & MacFadden, B.J. (Eds).
3 Cambridge: Cambridge University Press.
- 4 Jarman, P.J. & Sinclair, A.R.E. (1979). Feeding strategy and
5 the pattern of resource-partitioning in ungulates. In *Seren-*
6 *gueti, dynamics of an ecosystem: 130–163.* Sinclair, A.R.E.
7 & Norton-Griffiths, M. (Eds). Chicago: University of
8 Chicago Press.
- 9 Kendall, D.G. (1986). Size and shape spaces for landmark
10 data in two dimensions: comment. *Stat. Sci.* **1**, 222–226.
- 11 Kleiber, M. (1932). Body size and metabolism. *Hilgardia* **6**,
12 315–353.
- 13 Koenker, R. & Bassett, G.J. (1978). Regression quantiles.
14 *Econometrica* **46**, 33–50.
- 15 Koenker, R. & Machado, J.A.F. (1999). Goodness of fit and
16 related inference processes for quantile regression. *J. Am.*
17 *Stat. Assoc.* **94**, 1296–1310.
- 18 Langer, P. (2002). The digestive tract and life history of small
19 mammals. *Mamm. Rev.* **32**, 107–131.
- 20 Meloro, C. & O’Higgins, P. (2011). Ecological adaptations of
21 mandibular form in Fissiped Carnivora. *J. Mamm. Evol.*
22 **18**, 185–200.
- 23 Mendoza, M., Janis, C.M. & Palmqvist, P. (2006). Estimating
24 the body mass of extinct ungulates: a study on the use of
25 multiple regression. *J. Zool. (Lond.)* **270**, 90–101.
- 26 Millien, V. & Bovy, H. (2010). When teeth and bones disa-
27 gree: body mass estimation of a giant extinct rodent. *J.*
28 *Mammal.* **91**, 11–18.
- 29 Milne, N. & O’Higgins, P. (2002). Inter-specific variation in
30 Macropus crania: form, function and phylogeny. *J. Zool.*
31 *(Lond.)* **256**, 523–535.
- 32 Myers, T.J. (2001). Prediction of marsupial body mass. *Aust.*
33 *J. Zool.* **49**, 99–118.
- 34 Owen-Smith, N. (1988). *Megaherbivores. The influence of very*
35 *large body size on ecology.* Cambridge: Cambridge Univer-
36 sity Press.
- 37 Pérez-Barbería, F.J. & Gordon, I.J. (2001). Relationships
38 between oral morphology and feeding style in the ungulata:
39 a phylogenetically controlled evaluation. *Proc. Roy. Soc.*
40 *Lond. Ser. B* **268**, 1023–1032.
- 41 Peters, R.H. (1983). *The ecological implications of body size.*
42 Cambridge: Cambridge University Press.
- 43 Reed, K.E. (1998). Using large mammal communities to
44 examine ecological and taxonomic structure and predict
45 vegetation in extant and extinct assemblages. *Paleobiology*
46 **24**, 384–408.
- 47 Reguero, M.A., Candela, A.M. & Cassini, G.H. (2010).
48 Hypsodonty and body size in rodent-like notoungulates.
49 In *The paleontology of Gran Barranca: evolution and envi-*
50 *ronmental change through the Middle Cenozoic of Patago-*
51 *nia: 358–367.* Madden, R.H., Carlini, A.A., Vucetich, M.G.
52 & Kay, R.F. (Eds). Cambridge: Cambridge University
53 Press.
- 54 Reguero, M.A., Dozo, M.T. & Cerdeño, E. (2007). A poorly
55 known rodentlike mammal (Pachyrhinae, Hegetotherii-
dae, Notoungulata) from the Deseadan (Late Oligocene) of
Argentina. Paleocology, biogeography, and radiation of
the rodentlike ungulates in South America. *J. Paleontol.* **81**,
1301–1307.
- Riggs, E.S. (1935). A skeleton of *Astrapotherium*. *Geol. Ser.*
Field Mus. Nat. Hist. **6**, 167–177.
- Savage, V.M., Gillooly, J.F., Woodruff, W.H., West, G.B.,
Allen, A.P., Enquist, B.J. & Brown, J.H. (2004). The pre-
dominance of quarter-power scaling in biology. *Funct. Ecol.*
18, 257–282.
- Scarano, A.C., Carlini, A.A. & Illius, A.W. (2011). Interther-
iidae (Tyotheria: Notoungulata), body size and paleoecol-
ogy characterization. *Mamm. Biol.* **76**, 109–114.
- Scott, K.M. (1990). Postcranial dimensions of ungulates as
predictors of body mass. In *Body size in mammalian paleo-*
biology: estimation and biological implications: 301–335.
Damuth, J. & MacFadden, B.J. (Eds). Cambridge: Cam-
bridge University Press.
- Scott, W.B. (1910). Mammalia of the Santa Cruz beds. Part I.
Litopterna. In *Reports of the Princeton University expedi-*
tions to Patagonia, 1896–1899: 156. Scott, W.B. (Ed.). Prin-
ceton: Princeton University.
- Scott, W.B. (1928). Mammalia of the Santa Cruz beds. Part
IV. Astrapotheria. In *Reports of the Princeton University*
expeditions to Patagonia, 1896–1899: 301. Scott, W.B. (Ed.).
Princeton: Princeton University.
- Shapiro, S.S. & Wilk, M.B. (1965). An analysis of variance
test for normality (complete samples). *Biometrika* **52**, 591–
611.
- Smith, R.J. (1984). Allometric scaling in comparative biology:
problems of concept and method. *Am. J. Physiol.* **246**,
R152–R160.
- Smith, R.J. (1993). Logarithmic transformation bias in allom-
etry. *Am. J. Phys. Anthropol.* **90**, 215–228.
- Snowdon, P. (1991). A ratio estimator for bias correction in
logarithmic regression. *Can. J. For. Res.* **21**, 720–724.
- Tauber, A.A. (1997). Biostratigraphy of the Santa Cruz For-
mation (Lower Miocene) in southeastern Patagonia. *Ameg-*
hiniana **34**, 413–426.
- Villafañe, A.L. (2005). *Paleoecología de los Protheroitheriidae*
(Mammalia: Litopterna): un estudio basado en los cambios
en la masa corporal. Lic. thesis, Universidad Nacional de
La Patagonia San Juan Bosco at Trelew.
- Vizcaino, S.F., Bargo, M.S., Kay, R.F., Fariña, R.A.,
Giacomo, M.D., Perry, J.M.G., Prevosti, F.J., Toledo, N.,
Cassini, G.H. & Fernicola, J.C. (2010). A baseline pale-
oecological study for the Santa Cruz Formation (late-early
Miocene) at the Atlantic coast of Patagonia, Argentina.
Palaeogeogr. Palaeoclimatol. Palaeoecol. **292**, 507–
519.
- Vizcaino, S.F., Bargo, M.S., Kay, R.F. & Milne, N. (2006).
The armadillos (Mammalia, Xenarthra, Dasypodidae) of
the Santa Cruz Formation (early-middle Miocene): an
approach to their paleobiology. *Palaeogeogr. Palaeoclima-*
tol. Palaeoecol. **237**, 255–269.

Body mass of Santacrucian ungulates

G. H. Cassini, S. F. Vizcaino and M. S. Bargo

1 Warton, D.I., Wright, I.J., Falster, D.S. & Westoby, M.
2 (2006). Bivariate line-fitting methods for allometry. *Biol.*
3 *Rev. Camb. Philos. Soc.* **81**, 259–291.
4 Zar, J.H. (1999). *Biostatistical analysis*. 4th edn. Upper Saddle
5 River: Prentice Hall.

7 **Supporting Information**

8 Additional Supporting Information may be found in the
9 online version of this article:

Appendix SI Species of Santacrucian ungulates analyzed
and their body mass estimates. 10
11

Appendix SII Extant ungulates reference sample. 12
13

Please note: Wiley-Blackwell is not responsible for the content
or functionality of any supporting materials supplied by the
authors. Any queries (other than missing material) should be
directed to the corresponding author for the article. 14
15
16
17

AUTHOR QUERY FORM

Dear Author,

During the preparation of your manuscript for publication, the questions listed below have arisen. Please attend to these matters and return this form with your proof.

Many thanks for your assistance.


Query References	Query	Remarks
1	WILEY-BLACKWELL: Please confirm if the article type is correct.	
2	AUTHOR: Please confirm the postal address and telephone number are correct for the corresponding author.	
3	WILEY-BLACKWELL: Please provide the editor's name for this paper.	
4	AUTHOR: Please confirm the affiliation addresses are correct.	
5	AUTHOR: Please update the year of publication for the reference Cassini <i>et al.</i> , in press.	
6	AUTHOR: Please update the year of publication for the reference Cassini & Vizcaíno, in press.	
7	AUTHOR: Online Appendix I and II have been renamed as Appendix SI and SII. Please confirm that this is correct.	
8	AUTHOR: Please give manufacturer information for Microscribe G2L digitizer: company name, town, state (if USA), and country.	
9	AUTHOR: 'Although working with fragmentary fossils. . . .' This sentence has been reworded for clarity. Please confirm if this is correct.	
10	AUTHOR: 'On the contrary, <i>bm</i> estimations. . . .' This sentence has been reworded for clarity. Please confirm if this is correct.	
11	AUTHOR: Jarman and Siclair 1979 has been changed to Jarman & Sinclair 1979 so that this citation matches the Reference List. Please confirm that this is correct.	
12	AUTHOR: Please update the year of publication for the reference Cassini & Vizcaíno, in press.	
13	AUTHOR: Please update the year of publication and page range for the reference Cassini <i>et al.</i> , in press.	
14	AUTHOR: Please update the year of publication, volume number and page range for the reference Cassini & Vizcaíno, in press.	
15	AUTHOR: Please update the volume number and page range for the reference Elissamburu, 2011.	
16	AUTHOR: Please verify if the journal title in the reference Pérez-Barbería & Gordon, 2001 is correct.	
17	AUTHOR: Please verify changes in the footnotes of Tables 4 and 6.	

USING e-ANNOTATION TOOLS FOR ELECTRONIC PROOF CORRECTION


Required software to e-Annotate PDFs: Adobe Acrobat Professional or Adobe Reader (version 8.0 or above). (Note that this document uses screenshots from Adobe Reader X)

The latest version of Acrobat Reader can be downloaded for free at: <http://get.adobe.com/reader/>

Once you have Acrobat Reader open on your computer, click on the [Comment](#) tab at the right of the toolbar:



This will open up a panel down the right side of the document. The majority of tools you will use for annotating your proof will be in the [Annotations](#) section, pictured opposite. We've picked out some of these tools below:



1. Replace (Ins) Tool – for replacing text.




Strikes a line through text and opens up a text box where replacement text can be entered.

How to use it

- Highlight a word or sentence.
- Click on the [Replace \(Ins\)](#) icon in the Annotations section.
- Type the replacement text into the blue box that appears.

standard framework for the analysis of microeconomics. Nevertheless, it also led to the emergence of a new paradigm of strategic behaviour. The number of competitors in the industry is that the structure of the industry is a key component of the main components of the industry. At the level, are expected to be important works on the industry by Shiraz (M henceforth) we open the 'black b



2. Strikethrough (Del) Tool – for deleting text.



Strikes a red line through text that is to be deleted.

How to use it

- Highlight a word or sentence.
- Click on the [Strikethrough \(Del\)](#) icon in the Annotations section.

there is no room for extra profits and the number of competitors are zero and the number of (net) values are not determined by Blanchard and ~~Kiyotaki~~ (1987), perfect competition in general equilibrium of aggregate demand and supply in a classical framework assuming monopoly. An exogenous number of firms

3. Add note to text Tool – for highlighting a section to be changed to bold or italic.



Highlights text in yellow and opens up a text box where comments can be entered.

How to use it

- Highlight the relevant section of text.
- Click on the [Add note to text](#) icon in the Annotations section.
- Type instruction on what should be changed regarding the text into the yellow box that appears.

dynamic responses of mark ups consistent with the **VAR** evidence

sation... y Ma... and... on n... to a... on... stent also with the demand-



4. Add sticky note Tool – for making notes at specific points in the text.




Marks a point in the proof where a comment needs to be highlighted.

How to use it

- Click on the [Add sticky note](#) icon in the Annotations section.
- Click at the point in the proof where the comment should be inserted.
- Type the comment into the yellow box that appears.

and supply shocks. Most of the... a... number... dard fr... cy. Nev... ole of st... ber of competitors and the imp... is that the structure of the secto



USING e-ANNOTATION TOOLS FOR ELECTRONIC PROOF CORRECTION


5. Attach File Tool – for inserting large amounts of text or replacement figures.



Inserts an icon linking to the attached file in the appropriate place in the text.

How to use it

- Click on the [Attach File](#) icon in the Annotations section.
- Click on the proof to where you'd like the attached file to be linked.
- Select the file to be attached from your computer or network.
- Select the colour and type of icon that will appear in the proof. Click OK.



6. Add stamp Tool – for approving a proof if no corrections are required.




Inserts a selected stamp onto an appropriate place in the proof.

How to use it

- Click on the [Add stamp](#) icon in the Annotations section.
- Select the stamp you want to use. (The [Approved](#) stamp is usually available directly in the menu that appears).
- Click on the proof where you'd like the stamp to appear. (Where a proof is to be approved as it is, this would normally be on the first page).

of the business cycle, starting with the
 on perfect competition, constant ret
 production. In this environment goods
 extra profits and the market for marke
 he market for goods is determined by the model. The New-Key
 otaki (1987), has introduced produc
 general equilibrium models with nomin
 and... Most of this literature

APPROVED




7. Drawing Markups Tools – for drawing shapes, lines and freeform annotations on proofs and commenting on these marks.

Allows shapes, lines and freeform annotations to be drawn on proofs and for comment to be made on these marks..

How to use it

- Click on one of the shapes in the [Drawing Markups](#) section.
- Click on the proof at the relevant point and draw the selected shape with the cursor.
- To add a comment to the drawn shape, move the cursor over the shape until an arrowhead appears.
- Double click on the shape and type any text in the red box that appears.



For further information on how to annotate proofs, click on the [Help](#) menu to reveal a list of further options:

